

We would like to thank the reviewer for careful reading of this manuscript and for the constructive comments. We have responded to all comments below, added explanations where necessary, and applied most of the suggested changes or corrections.

We have updated our analysis with new present climate runs, which were not available previously. Unlike the historical 1979-2005 simulation, the new runs now follow the HAPPI experiment protocol, i.e. they constitute an ensemble of decadal runs in 2006-2015. Therefore they are more suitable for the analysis investigating changes between the present and future climate. With the updated/improved design of the analysis, some of the reviewer's comments may not apply to the current version of the paper. We will provide more explanations below.

Anonymous Referee #1

Received and published: 17 December 2017

1 General Comments

The manuscript "Euro-Atlantic winter storminess and precipitation extremes under 1.5 °C versus 2 °C warming scenarios" by Barcikowska et al presents comparisons between storminess and precipitation in the 20th century and in the early 22nd century using newly available model simulations from the HAPPI project in different horizontal resolution. They first evaluate model results through comparing results from the model runs on different horizontal resolutions with ERA-Interim (circulation-type variables). Here, they conclude that the 0.25 degree resolution provides the best results, where atmospheric features are presented superior to the lower horizontal resolution model simulations. In the following, 0.25 degree model precipitation is compared with data from the EObs and GHCN datasets, where the authors find very good agreement.

Afterwards, the authors investigate the differences between the scenarios under 1.5 and 2 °C warming, whereby they present changes in the mean-state of the large-scale atmospheric circulation and precipitation, in daily and sub-daily precipitation and wind extremes, and in storminess for the 0.25 degree run. Overall, the manuscript deals with an important subject and combines different aspects of how storminess and precipitation changes under 1.5 and 2 °C warming, also with regard to making the model simulation finer. The manuscript clearly conveys this subject, but nevertheless suffers from several major aspects that need improving and/or further clarification, before it is ready to be published.

1. The HAPPI ensemble consists of several model runs from different modeling centers (not mentioned in the text). I understand that you concentrate on the CAM5-simulations, but give no reason why the other simulations are discarded.

AU: In this study, we only focus on the same model version run at different resolutions. This allows us to investigate the impacts of a very high model resolution on the representation of large-scale and regional features in comparison to a coarser resolution. Additionally CAM5 1.2-0.25 provides unprecedented opportunity to investigate extremes on subdaily time scales. We stated this in the introduction, but will make it clearer in the revised version. Also we don't feel obliged to list all the modeling centers participating in the HAPPI projects. In similar way, studies employing particular CMIP models don't usually list all of the other centers/models, contributing to the CMIP models community. (page 5, line 30-35)

2. Regarding the model resolution: The implications found for the large-scale atmospheric circulation cannot be overstated enough and put the model into the sphere of dynamically downscaled regional models with corrections for the larger scales. I agree with section 3.4 that there is room for some kind of sensitivity study here.

AU: We agree but think that such sensitivity studies should be part of a separate study. We nevertheless motivate such studies in our manuscript.

3. I also find it very interesting to see the differences between the present climate and 1.5 °C vs 2 °C warming. Your results suggest (as you wrote) that there seems to be a threshold in between, that once crossed, exacerbate storminess conditions.

AU: Yes, one interpretation could be associated with the threshold. Other interpretation, with an additional analysis including new present climate simulations, suggests that the difference might be due to the asymmetry in aerosol forcing between the present and future climate. It is a very important point and we elaborate on that in the revised version. We support our statements, by updating the analysis with new simulations of the present climate. These simulations follow the HAPPI protocol (unlike the previous ones) and are more relevant to address this issue. **(page 7, lines 3-9; 27-29; page 11, lines: 1-9, page 12, lines: 10-21; page 17, lines 25-30)**

4. Why is ERA Interim the reference for midlatitude atmospheric circulation? How does ERA Interim compare to other reanalyses with regard to circulation?

AU: We use ERA interim, because its spatial resolution is comparable to the high-resolution CAM5 model simulations. ERA interim has considerably higher model resolution (80 km at 60 vertical levels) than other reanalysis products. Hence it is able to resolve sharper spatial gradients than e.g. NCEP/NCAR Reanalysis. Hodges et al. 2011 compares reanalyses for extratropical cyclones and shows that the newer reanalyses (especially ERA-Interim and NCEP-CFSR) agree (both in terms of numbers and locations) much better than the older ones (JRA-25) for both hemispheres and that intensities are higher. As our purpose is to validate the model, ERA interim serves the purpose better than the coarse reanalysis products like NCEP/NCAR.

We updated the Figure 1a and relevant text in the manuscript, by adding the comparison of the ERA-I and NCEP-DOE data sets. It shows that differences between the observations are much smaller than those derived between observations and CAM5. See changes in 2.1 section: **page 7 line,14-16; page 9 line: 13-22.**

The figures attached below differentiates top) ERA-I and CFSR, bottom) NCEP and ERA-I, showing that for the North Atlantic-European region, the largest observational uncertainty is located over Greenland. Observational uncertainty is much smaller, compared to the differences between the observations and the model.

We also added reference (page 9, line 16):

Hodges, K.I., R.W. Lee, and L. Bengtsson, 2011: A Comparison of Extratropical Cyclones in Recent Reanalyses ERA-Interim, NASA MERRA, NCEP CFSR, and JRA-25. *J. Climate*, 24, 4888–4906, <https://doi.org/10.1175/2011JCLI4097.1>




Figure. Time-mean average of the DJF sea level pressure [hPa] over the period 1979-2005, re-gridded to the  $2.5^{\circ} \times 2.5^{\circ}$  horizontal grid for top) ERA-Interim (shaded) and differences relative to CFSR (at original resolution T382,  $\sim 0.34^{\circ}$  resolution at equator); bottom) NCEP/DOE 2 ( $\sim 2.5^{\circ}$  original resolution, shaded). Contours show differences relative to ERA-Interim (ERA-I,  $\sim 0.75^{\circ}$  lat-lon original resolution)

5. The changes of the SLP gradient are interesting. As you mention the NAO in the beginning of section 3, what are the consequences for the NAO index caused by the increasing SLP gradient difference? You could compute the NAO index and show how it changes as it should be a stationary process centered around 0 in the long-term.

AU: Because of the strengthening of the SLP gradient, the NAO index will likely have a positive tendency, when compared to a +2C future with the +1.5C/present climate experiments. However under each stabilization scenario, the spatial pattern/definition of the NAO would be different. Hence in this experimental setup, the NAO could be rather investigated separately for each of the three ensembles. Therefore a more thorough analysis could be done only on the NAO, but in a separate study and based on transient simulations rather than time slice experiments.

6. This one is very important: The question of statistical significance has not been dealt with properly and currently is rather imprecisely given (sections 2.2, 3.2 and S4). S4 and p.12 118 ("which defies statistical significance"): What you actually show is the distribution of differences, from which you can infer a confidence interval. What you do not get are real implications about statistical significance as written.

Please repeat your analysis here. Also: please move the detailed description of your method into the method-section without repeating the details later. Regarding S4: The figure looks very choppy. Either it is showing some kind of histogram sampled for specific blocks of differences (then it should be stated clearly), or it demonstrates undersampling in your bootstrapping approach. Either way, it would be good to redo the bootstrapping with a bigger sample than just 1000. The computation is cheap and very likely results in a better representation of the distribution of differences.

AU: This is indeed an important point. We would certainly repeat this analysis, at the absence of present climate simulations following the HAPPI protocol. Instead, we will update the analysis with new simulations and discussion. These are tailored specifically to remove the effects of different phase of internal variability, which was questioning the statistical significance of the derived results.

In the new version of section 3.1, we were now able to use new ensemble simulations of the present climate. These runs represent the 2006-2015 period, instead the previously used simulation of the 1979-2005 period. The new runs follow closely HAPPI protocol, and thus facilitate more direct interpretation of the results. The new present climate runs, as well as each of the future warming, i.e. +1.5C and +2C experiment includes internal climate SST variations (e.g. ENSO) during the same decadal period, i.e. 2006-2015. Therefore it is expected that the impacts of internal variations will be canceled out while discriminating between all three experiments. In this context we found the bootstrapping approach to be redundant and we removed this part of analysis (and associated figure S4).

Please note that for the analysis of the climatological features of the model and the impact of the resolution (e.g. section 3, and 3.3), we used historical runs for 1979-2005.

It is worth noting, that the asymmetry in forcing, when comparing the present and future climate experiments, remains a factor complicating the interpretation of the results. The changes associated with warming at the 1.5°C level stem from an interplay of a number of forcings, including strong aerosol reductions, while an additional half a degree warming is solely a consequence of further CO<sub>2</sub> increase and ocean warming (**page 17, line 25-30**). Please see more explanation: page 11, lines: 1-10, page 12, line: 15-20.

7. The meridional SLP gradient and its differences: Sections 3 and 3.2 write about the SLP gradient, but only refer to figures 4 and 5 showing the respective MSLP plots. I, as a reader, am not able to estimate the gradient and gradient differences from such plots. As you define the gradient in section 2.2, it is very difficult to relate the Azores-Icelandic pressure difference to plots of MSLP or MSLP differences, even though I know about the related atmospheric patterns. Why not just give the gradient as a number somewhere? (also for 1.5 °C and 2 °C scenarios, and the differences).

AU: The estimated changes in the SLP gradient are consistent with the derived changes in the large-scale circulation. We included the numbers and the relevant discussion in the text: section 3.1 **page 11, line: 15-18, page 12: 12-14, 15-20.**

We also underlined the fact that the estimations of the differences between the +1.5C and +2°C scenario are most pronounced and that we will focus mostly on the differences derived between the +1.5C and +2°C scenario. The explanation is provided in the page **12, line 15-21.**

Additionally, contours (labeled) of SLP in Figure 4a and 5a show the time-average SLP in the present climate, while the differences between the experiments are shaded. This facilitates the interpretation of the future SLP changes in the context of the mean ambient flow.

Another note to the SLP gradient: It suddenly appears at the end of section 2.2 without prior mentioning. It should be introduced a little earlier along with the other variables (p6 l 32ff) including the reason to do so.

AU: Yes, we provided and expanded the information on the SLP gradient in the earlier part of the methods section (**page 7, line 37 - page 8 line 1-4**).

8. For section 3, can you provide spatial statistics, such as the pattern correlation when you describe the resemblance of simulations with observational datasets?

AU: Yes, we have updated the analysis with the pattern correlation.

The observed average SLP patterns, derived from ERA-I and NCEP/DOE 2 (~2.5°) share a correlation (uncentered pattern correlation) of 0.98. The differences between the ERA-I and CAM5 are larger, than the observational differences.

The magnitude and the pattern in ERA-I correlate best with the one simulated at similar horizontal resolution (CAM5\_1), returning number of 0.96. Correlations with the remaining two are slightly smaller, i.e 0.95 for CAM5\_2 and 0.94 for CAM5\_0.25. Please see in text (**Section 3, page 9: lines 26-29**).

9. Section 3.2, p. 12, l 19 "time-average over 1979-2005": Do you take care of any secular trend, which might be imminent in 26 years of data, but may disappear in a shorter time period?

AU: That is a valid point, which we would certainly take into consideration. However, in the updated version of the analysis in this section, we no longer use the 1979-2005 run. Instead, we use new runs of the present climate. These simulations are more suitable as they are set to follow the HAPPI protocol. We will update the revised version of the paper accordingly.

In this version we are using decadal runs (2006-2015). The impact of secular trend in these runs is rather negligible. (methods section: **page 7, line 28-29**)

3. Technical comments:

Some passages of the manuscript are not concise (e.g. repetition of methods in the text, when such details belong to the method section). Sometimes the manuscript does not read well.

AU: We worked on readability of the manuscript, i.e. section 3 is shortened and clarified. We merged section 3.1 and 3.2. Some technical information is moved to the method section (**page 7, line 28-29, page 7, line 36-37; page 8, line 1-3**). But we insist on repeating some technical facts in the introduction of section 3.1, instead mentioning them only in the methods, to facilitate the interpretation of the following results. (**page 10, line 35 - page 11, line 10**).

1. Please check the references. There are references clearly missing, for instance Barcikowska et

al., 2017 or Gilleland and Katz, 2014; or misleading like Feser et al., 2014 (did you mean Feser et al., 2015?). There might be more that I have overlooked.

AU: We corrected the references in the revised version.

2. Section 3.1 feels a little superfluous and could easily be merged into the method section.

AU: The section will be shorter and more concise in our revised version, as it excludes the analysis of internal variability impacts.

As suggested, we merged section 3.1 and old version of section 3.2. As mentioned above, some part of information, e.g. description of the analysis of the large-scale circulation is moved to the method section. However we left the information, which facilitates the interpretation of the following results. (page 10, line 35-37; page 11: line:1- 9)

AU: We have applied corrections, as suggested below:

3. p 11 32: the British Isles → corrected (page 4, line 13)

4. p 41 21: Zappa et al. (2014) have shown: → corrected (page 4, line 22)

5. p 61 4: provided by the C20C+ Detection and Attribution Project → corrected

6. p 61 7: the CAM5-1-1degree run [...] and the CAM5-1-0.25degree run (missing articles) → corrected (page 6, line 13-15)

7. p 61 9: add an "and" before 0.3125\_x0.234 → corrected (page 6, line 16)

8. p 61 10: remove the last ")" → corrected (page 6, line 18)

9. p 61 18: do you need commas in front and after the subclause "using atmosphereonly models?" → we think that commas will improve clarity of the sentence (page 6, line 25)

10. p 61 22: remove comma after offset → corrected (page 6, line 29)

11. p 61 35: zonal wind; what about meridional winds or wind speeds in general? → edited as "winds" (page 7, line 11)

12. p 71 17: Wilcoxon signed rank test, can you add a reason why you use it? → added in page 7: line 31-32

13. p 71 21: a block (seasonal) maximum → corrected (page 8, line: 9)

14. p 71 22: The whole sentence with "Assumptions that our analyzed data..." needs rewriting. → corrected (page 8, line 9-10)

15. p 71 26: there is something wrong with (1-1/T)th (you accidentally inserted a comma) → corrected (page 8, line 12)

16. p 71 33-35: please add a reference → corrected (page 8, line: 23)

17. p 91 24: will presumably lead to a → corrected (page 10, line 9)

18. p 91 37: either from the model bias or from observational bias → corrected (page 10, line 22)

19. p 101 4: high-resolution runs provide a more accurate representation → corrected (page 10, line 26)

20. p 101 20: provided for the years → we removed this sentence, since we no longer use the 1979-2005 simulation in the section 3.1

21. p 101 20: internal SST variations being in a different phase → the same as above

22. p 111 18-21: reflect, generally after reductions, CO2 increase

23. Section 3.4: This one reads very well (also applies to section 2.3).

24. Figure 7: What is a fractional difference? → We corrected it as a fractional change (difference in a relation to the mean in a reference period. (please see Fig 7 caption and text in page 13, line 10)

25. Figure 9: The caption talks about 50 years. As far as I understood the manuscript, shouldn't it be less years? Maybe I did not get, where the 50 years sample comes from? → The sample comes from the 5-member ensemble of decadal runs. We've clarified it in the revised manuscript (Figure

9 caption).

26. Figure S2: There is either something wrong with the figure caption or there is a whole figure missing. → We removed figure S2. The figure is not relevant for the analysis (section 3.1) in the current version, as it employs the 1979-2005 runs (instead 2006-2015). Because of the same reason we removed the old figure S3, which was showing the differences between the future and the 1996-2005 runs.

We would like to thank the reviewer for careful reading of this manuscript and for the constructive comments. We have responded to all comments below, added explanations where necessary, and applied most of the suggested changes or corrections.

We have also updated our analysis with new present climate runs, which were not available previously. Unlike the historical 1979-2005 simulation, the new runs now follow the HAPPI experiment protocol, i.e. they constitute an ensemble of decadal runs in 2006-2015. Therefore they are more suitable for the analysis investigating changes between the present and future climate. This approach is also more suitable to address the main reviewers concerns on the non-linear effects of the derived climate change, shown in the study. We will provide additional explanations below.

Anonymous Referee #2

A: Based on their analysis of the high resolution time slice experiments, the authors state that most of the changes due to anthropogenic forcing up to the 2C level only start occurring after the 1,5C threshold has been surpassed, and claim this may be due to non-linearities in the climate system (page 17, around line 30). While this may well be true to some point, I find it very hard to believe that while the climate change signal e.g. DJF MSLP and 850 hPa winds up to a 1,5C warmer world is practically zero (cf. Fig. S1a), we become a very strong response when we add the additional 0,5C (cf. Fig. S1b). The same could be argued for the other fields.

In my view, such a strong difference for such a small increment of external forcing could only come from (a) internal variability (for example, storminess is at a low level of its decadal variability in the period chosen for the 1,5C experiment, while it is on a high level on the period chosen for 2C) and/or (b) there are some issue in the set up of the high resolution simulations which have lead to these differences, and not the small change in the forcing. This does not turn the results per themselves wrong, but means that the authors may be misinterpreting (or at least over-interpreting) their results. It can be that the problem is related with the SST set up, as the authors shortly discuss in section 2.1. For example, how does the climate change signal look like for the (hopefully transient) lower resolution GCM simulations? How does the climate change signal look like for the single ensemble members? This would be important to analyse in detail to identify if the changes in precipitation and wind are continuous over time (plus natural decadal climate variability) or if indeed some strong “non-linear” effects occur. It is very improbable that the AMOC will collapse in a two degree warmer world. What might be important here is the thermodynamical effect (Clausius-Clapeyron), primarily for precipitation intensity. This is very important to rule out that the obtained results are not simply caused by some issue with the set up of the model simulations and clearly relate the changes to the increases of anthropogenic forcing.

AU: Thank you for the comment. Indeed, the “non-linear” effects of the derived changes, when comparing first half a degree warming (+1.5C minus present) with the additional half a degree warming (+2C minus +1.5C), are likely associated with the simulations set up. i.e. an asymmetry in the aerosol forcing between the present and future climate scenarios. The changes associated with warming at the 1.5°C level stem from an interplay of a number of forcings, including strong aerosol reductions, while an additional half a degree warming is solely a consequence of CO<sub>2</sub> increases and ocean warming.

We enhanced the interpretation in the new revised version:

Section 2.1: page 7, lines: 2-7

Section 3.1: page 10, lines: 35-37; page 11, lines: 1-9; page 12, lines: 10-21

Section 4: Page 17, lines 31-37

Minor comments:

#1 Page 2, line 34: regarding the role of cyclone clustering for the wind and flood impacts, the authors could refer for example to Priestley et al. (2017), which deals specifically with this topic.

AU: Thank you, we included the reference (Page 2, line 33)

#2: Page 3, around line 10: The “poleward” shift of the storm tracks due to enhanced anthropogenic forcing is correct on zonal average, but this is not always true for Europe: while this is clear for the summer half year, the results for the winter, particularly DJF (the focus here), are different – see your own text in page 4, line 10. Please check e.g., the discussion in Zappa et al (2015), and also their Fig.1.

AU: Yes, we agree that the “poleward shift” is a simplified statement. But it is a valid statement in the used context. We also added, that more discussion on the spatial changes of storms in different seasons will be included in latter part. (Page 3, line 12-13)

#3: Page 4, line 12: The results by Zappa et al. (2013b) do not contradict the results described in many of the papers mentioned in the previous sentence. For example, the results from Bengtsson et al. (2006), Pinto et al. (2009) or Zappa et al. (2013b) only show small differences in detail regarding the intensification of cyclone activity over the British Isles and the strong decrease in the Mediterranean. Thus, “however” is not the correct word here, maybe “Moreover” would be more appropriate.

AU: Thank you, we applied the correction. (Page 4, line 13)

#4: Page 4/5: The authors describe their reasoning as if it would be the very first time that high-resolution global climate modelling is performed at 25km resolution. This is not correct – for example, there is a whole EU project regarding high resolution modelling to study impacts of climate change for Europe (PRIMAVERA, <https://www.primavera-h2020.eu/about/objectives/>). It is strange not to mention this at all in the introduction, nor any of the related papers (e.g. Schiemann et al., 2017). The MIROC and MRI groups has also been working high-resolution climate models for several years (e.g. Murakami et al., 2011), which is not really discussed– though Kitoh and Endo (2016) is shortly mentioned. Please note that another possibility would be to use RCMs as ESMs (e.g. Sein et al., 2015). Please enhance.

AU: Thank you for the correction, we enhanced the Introduction. We included the information about PRIMAVERA project and references (Schiemann et al., 2017, Sein et al., 2015). As the scope of the section is to provide an introduction on extratropical storms, we skipped one reference (Murakami et al. 2011), which deals with tropical cyclones. Instead we refer to Yang et al. 2015.

Yang, X., G.A. Vecchi, R.G. Gudgel, T.L. Delworth, S. Zhang, A. Rosati, L. Jia, W.F. Stern, A.T. Wittenberg, S. Kapnick, R. Msadek, S.D. Underwood, F. Zeng, W. Anderson, and V. Balaji, 2015: Seasonal Predictability of Extratropical Storm Tracks in GFDL’s High-Resolution Climate Prediction Model. *J. Climate*, 28, 3592–3611, <https://doi.org/10.1175/JCLI-D-14-00517.1>

#5: References: many issues, both in the reference list and in the text. For example, Zappa et al (2012) is surely one of the two Zappa et al. (2013) papers, Feser et al (2014) is probably Feser et al. (2015). Pinto et al (2009) is three times in the reference list, and several references mentioned in the text are not included (e.g. Barcikowska et al. 2017) are not included in the Reference List. Please enhance.

AU: We corrected and updated the citations and the references list.

#6: Figures: it would be very good to had labels to the isolines in the figures, particularly in Figures 8 and 9. It is not enough to mention the values in the captions. Please enhance.

AU: The figures are now updated and include isolines in the Figure 8. The corresponding figure captions are also updated.

# Euro-Atlantic winter storminess and precipitation extremes under 1.5°C versus 2°C warming scenarios

Monika J. Barcikowska<sup>1</sup>, Scott J. Weaver<sup>2</sup>, Frauke Feser<sup>3</sup>, Simone Russo<sup>4</sup>, Frederik Schenk<sup>5</sup>, Dáithí A. Stone<sup>6,7</sup>, **Michael F. Wehner**<sup>6</sup>, Matthias Zahn<sup>3</sup>

5 <sup>1</sup>Environmental Defense Fund, New York City, United States

<sup>2</sup>Environmental Defense Fund, Washington, D.C, United States

<sup>3</sup>Institute of Coastal Research, Helmholtz Centre Geesthacht, Geesthacht, Germany

<sup>4</sup>European Commission, Joint Research Centre, Via Enrico Fermi, Ispra, Italy

10 <sup>5</sup>Bolin Centre for Climate Research, Department of Geological Sciences, Stockholm University, Stockholm, Sweden

<sup>6</sup>Lawrence Berkeley National Laboratory, Berkeley, CA, USA;

<sup>7</sup>Global Climate Adaptation Partnership, U.K.

*Correspondence to:* Monika J. Barcikowska (mbarcikowska@edf.org)

**Abstract.** Severe winter storms in combination with precipitation extremes pose a serious threat to Europe. Located at the south-east exit of the North Atlantic's storm track, European coastlines are directly exposed to impacts by high wind speeds, storm floods and coastal erosion. In this study we analyze potential changes in simulated winter storminess and extreme precipitation which may occur under 1.5°C or 2°C warming scenarios. Here we focus on a first simulation suite of the atmospheric model CAM5 performed within the HAPPI project and evaluate how changes of the horizontal model resolution impact the results regarding atmospheric pressure, storm tracks, wind speed and precipitation extremes.

20 The comparison of CAM5 simulations with different resolution indicates that an increased horizontal resolution to 0.25° is not only refining regional-scale information, but also improves large-scale atmospheric circulation features over the Euro-Atlantic region. The zonal bias in SLP and wind fields, which is typically found in low-resolution models, is considerably reduced. This allows us to analyze potential changes in regional- to local-scale extreme wind speeds and precipitation in a more realistic way.

25 Our analysis of the future response for the 2°C warming scenario generally confirms previous model simulations suggesting a poleward shift and intensification of the meridional circulation in the Euro-Atlantic region. Additional analysis suggests that this shift occurs mainly after exceeding the 1.5°C global warming level, when the midlatitude jetstream manifests a strengthening north-eastward. At the same time, this north-east shift of the storm tracks allows an intensification and north-east expansion of the Azores high leading to a tendency of less precipitation across the Bay of Biscay and North Sea.

30 Regions impacted by the strengthening of the midlatitude jet, such as the northwest coasts of the British Isles, Scandinavia and the Norwegian Sea, and over the North Atlantic east from Newfoundland experience an increase in the mean as well as daily and sub daily precipitation and wind extremes and storminess

suggesting an important influence of increasing storm activity in these regions in response to global warming.

## 1. Introduction

5 International climate policy discussions use annual mean globally averaged temperature targets as the metric to anchor climate mitigation and adaptation strategies. While useful for climate policy development and implementation, global temperature targets do not explicitly convey the climate impacts that may be felt by society at seasonal and regional scales and hence make it difficult to justify any target as a safe level of warming (Knutti et al. 2015). The recent Paris agreement (*Adoption of the Paris Agreement* FCCC/CP/2015/L.9/Rev.1, UNFCCC 2015) hopes to limit the rise in post-industrial globally averaged  
10 temperature to no more than 2°C, while pursuing efforts toward the more ambitious 1.5°C target. Accordingly, understanding the changes in regional climate as the result of this half a degree difference in these two global temperature levels is important to clarify projected near-term climate change impacts.

In this study, we focus on projected changes in winter storminess and precipitation extremes over the Euro-Atlantic region. The winter climate in the North-Atlantic-European sector is dominated by variations in  
15 mid-latitude westerly winds which determine the position and intensity of storm tracks and thus the pathways of momentum, moisture and temperature transport. Extra-tropical cyclones dominate the redistribution of energy with a net poleward heat transport. They typically form in the region of strong baroclinic activity at the (sub-)polar front of Arctic vs. (sub-)tropical air masses. Stronger pressure gradients are linked to increased storminess and precipitation over central and Northern Europe and less  
20 storms and precipitation over southern Europe and vice versa for weak pressure gradients (e.g. Pinto et al. 2009). Large-scale storminess is dominated by multi-decadal variations in response to a complex interplay of different factors which may lead to changes in storm track position and intensity. The location of the storm track generally changes seasonally in response to solar insolation. Here, changes in the position of the sea-ice front push storm tracks southward while tropical sea-surface temperatures build a barrier in the  
25 south (Shaw et al. 2016).

Owing to its exposure to the direct impact by cyclones from the North Atlantic, weather extremes in this region frequently cause profound socio-economic costs. Heavy rainfall and intense winds are often associated with extratropical cyclones, and may cause flooding and storm surge, damaging infrastructure, industry, agriculture, and forestry. As an example for the North Sea region, extreme wind gusts can exceed  
30 Cat. 3 Hurricane wind forces like during storm Christian/Allan on 28<sup>th</sup>/29<sup>th</sup> of October 2013 with 171 km/h at the German North Sea Coast and 193 km/h over Denmark (von Storch et al. 2014). Hydrological extremes like the coastal as well as inland flooding over the southern United Kingdom during winter 2013/14 (Schaller et al. 2016, **Priestley et al. 2017**) are also closely tied to unusual series of low pressure systems including severe storm clusters and persistent rain. Given the large spatial variation in winter  
35 European climate affected by Euro-Atlantic storminess, any effect of global climate change on storminess could profoundly contribute to the associated regional impacts.

Many observational studies on the hydrological cycle in the recent century show wettening tendencies in the northern hemisphere highlighted by annual precipitation increases over large portions of the European continent including Scandinavia and central–eastern Europe. While these tendencies have also been detected in the winter season over most of these regions, they are not present over the southern flanks (Maraun, 2013) leading to a north-south dipole structure in precipitation anomalies over the European sector. A similar dipole pattern, with positive sign tendencies for the north and negative sign for the south of the continent, were also found in the records of winter extreme rainfall (Donat et al., 2013, Fischer et al., 2014) and river flows (Stahl et al., 2010, 2012). Other studies (Casanueva et al., 2014; Fleig et al., 2015) linked these changes directly to the altered large–scale circulation patterns. Hov et al. (2013) have shown that the intensification of the winter heavy rainfall in northern and north-eastern Europe is directly associated with the observed poleward shift of the North Atlantic storm track and weakening of Mediterranean storms. **Nevertheless, spatial changes of the storm track activity in this region feature much higher complexity, as will be discussed in the latter part.**

There is however an insufficient understanding of long-term changes in storminess and their drivers (Seneviratne et al., 2012). Records of extreme winds suffer from large inhomogeneities, contributing to uncertainty of the derived statistics during period of the satellite era (Hartmann et al., 2013; Feser et al., 2015a) in addition to spurious long-term trends in global reanalysis data (e.g. Krueger et al., 2013; Schenk and Stendel, 2016). There is however consistency across multiple datasets and medium confidence in a poleward shift of storm tracks since the 2<sup>nd</sup> half of the 20<sup>th</sup> century (Seneviratne et al., 2012). The observed increase in northern hemispheric storminess towards northern latitudes and a decrease southwards during the past several decades is consistent with the northward shift of storm tracks and their intensity since at least 1970 (e.g. Ulbrich et al., 2009; Lehmann et al., 2011; Hov et al., 2013; Feser et al. 2015a). Wang et al. (2009) attributes these changes since 1950 at least partly to external drivers.

Recent efforts to better understand future impacts of global warming on the Euro-Atlantic climate and weather and their extremes such as mid-latitude storminess typically involve an assessment of changes to various properties of atmospheric dynamics in global climate models (e.g., changes in wind and sea-level pressure variance) under various Representative Concentration Pathway (RCP) greenhouse gas forcing scenarios (Yin, 2005; Lu et al., 2007; Wu et al., 2011; Feser et al., 2015b).

Projections of future annual precipitation indicate an increase for the northern parts and a decrease for the southern parts of Europe. Both global climate model-based (Sillmann et al., 2013; Giorgi et al., 2014) as well as regional climate model-based studies (Rajczak et al., 2013; Jacob et al., 2014) agree that the strongest increase in the winter heavy rainfall will occur over Scandinavia and eastern Europe. Moreover, Sillmann et al. (2013) have shown that heavy rainfall is projected to increase even in the regions with a mean precipitation decrease (e.g. over the Mediterranean region). Studies analyzing high-resolution, single-model projections (Kitoh and Endo, 2016; Barcikowska et al. 2017) corroborate these results. This bipolar pattern, with positive tendencies over the northern flanks of central and western Europe and a decrease over southern parts of Europe, has also been found in a multi-model ensemble projection (Donat et al. 2011) for

wind speeds.

Projections of future changes in the mid-latitude storms in the Northern Hemisphere indicate remarkable changes, however their features (e.g. spatial patterns and intensity) show a strong dependency on the analysis method as well as the generation of the models. Projections based on the ensemble mean of 16 CMIP3 (early 2000s generation) GCMs (Lambert and Fyfe 2006) as well as earlier modeling studies (Lambert, 1995, 2004) suggest a reduced frequency of extratropical cyclones due to a decreased surface meridional temperature gradient over the northern hemisphere. However, this decrease is not spatially uniform as storm activity south of 60° N over the northeast Atlantic and western Europe opposes this tendency, showing an increase in the CMIP3 projections (Leckebusch et al. 2006). Most of CMIP3 and earlier studies (Della-Marta and Pinto, 2009; Pinto et al., 2009b, Bengtsson et al., 2006; Geng and Sugi, 2003; Leckebusch et al., 2006; Pinto et al., 2006) indicate an eastward extension of storminess associated with an increase in frequency of strong storms over the British Isles, the North Sea and north-western Europe. **Moreover**, Zappa et al. (2013) has shown that the winter storm track's response in CMIP5 (late 2000s generation) projections manifests as a tripolar pattern, with an increase over the British Isles and decreased activity over both, the Norwegian and the Mediterranean Seas.

In most of the modeling applications, the horizontal resolution constrains the ability of GCMs to simulate both the important regional features and the large-scale circulation. So far, the quality of the simulated present climate and thus presumably projections of future climate have improved over time owing to progressing development of GCM's including resolution and representation of the physical process. Nevertheless, present climate simulations in CMIP5 models still suffer from notable biases i.e. on regional scale.

**Zappa et al. (2012) have shown** that CMIP5-based cyclones are generally too weak and the DJF storm track pattern is too zonal. These deficiencies are associated with the tripolar bias, manifested by negative anomalies over the Norwegian Sea and central-eastern parts of the Mediterranean, and positive anomalies spreading across north-western to central Europe towards the Black Sea. These biases are largely due to the inability of low-resolution models to correctly capture flow-orography interactions and thus correctly represent the tilt of the eddy-driven jet stream over the North Atlantic. Kelley et al. (2011) showed that the increased horizontal resolution in CMIP5 (~200 km) models potentially allowed for a spatial refinement in the simulated geographical pattern and for improvements in the simulated amplitude of precipitation indices. However the resolutions of the CMIP5 GCMs are not sufficiently high to correctly represent daily precipitation extremes (and their changes) and lead to severe underestimations (Sillmann et al., 2013; Van der Wiel et al., 2016).

Projections downscaled with Regional Climate Models (RCMs) may refine spatial details but will mostly inherit the large-scale circulation biases from the driving GCMs. Therefore, increasing spatial and temporal resolution in GCMs is crucial to **improve the representation of the simulated mean climate, weather extremes, and their changes. The PRIMAVERA project (<https://www.primavera-h2020.eu/about/objectives/>) focuses specifically on high-resolution modeling of the Euro-Atlantic**

climate. Modeling efforts pursued within this project facilitate an analysis of regional changes and associated impacts. For example Schiemann et al., 2017 have shown an improved representation of atmospheric blocking, which often redirects storm tracks, when simulated at higher (i.e. 25 km) resolution. Yang e al. 2015 used a high-resolution climate prediction model and highlight the importance of credibly resolved upper tropospheric jet flow, in order to skillfully predict storm-track statistics and associated extremes. Other studies (Kitoh and Endo, 2016; Barcikowska et al., 2017) employing relatively high-resolution models (~20 to ~50km) pointed to much higher skill in capturing both large-scale circulation features, spatial features and magnitude of precipitation extremes. First experimental simulations at even higher resolution (1-5 km, Kendon et al., 2014; Ban et al., 2015; Lehmann et al., 2015) were capable of projecting changes in heavy rainfall on sub-daily time scales but are usually too expensive to perform.

While it is important to understand the impacts from the worst case emissions scenarios, in order to support policy relevant mitigation and adaptation strategies as expressed in the Paris agreement, it is also necessary to assess the role of near-term global climate change in anticipating the shifts in regional climate and weather as a function of the 1.5°C and 2°C climate policy goals. However, there is a wide range of global temperature responses and considerable overlap of the CMIP5 models to lower emission scenarios that encompass the 1.5°C and 2°C levels of global warming (Mitchell et al., 2017). As such, teasing out the relative differences between these two temperature targets is not trivial and requires an alternate modeling strategy that obviates the transient uncertainty with respect to when a given model crosses either the 1.5°C or 2°C threshold (Kalmarkar and Bradley, 2017), mitigates the impact of potential differences in the phasing and amplitude of internal climate variability, and provides enough ensemble members to adequately distinguish the relevant climate change statistics.

The high resolution CAM5 simulations as part of the Half a Degree Additional warming, Prognosis, and Projected Impacts (HAPPI) project provides such a set of model experiments targeted specifically at differentiating the climate response between the 1.5°C and 2°C global temperature levels and their regional implications (Mitchell et al., 2017). The high spatiotemporal resolution of the CAM5 HAPPI experiments are unique in that they allow for a detailed analysis of large-scale changes to North Atlantic storm track activity and differential impacts as a function of model resolution - a necessary component for studying changes in precipitation and atmospheric circulation on sub-daily time-sales and for the representation of extreme weather events.

The aim of this study is to assess changes in the winter climate and weather extremes over the Euro-Atlantic region, associated with the 1.5°C and 2°C levels of global warming. In this study we employ the Community Atmospheric Model version 5 (CAM5), which is available at different horizontal resolutions. This allows us to investigate the impacts of a very high model resolution on the representation of large-scale and regional features in comparison to a coarser resolution. Additionally this model provides unprecedented opportunity to investigate extremes on sub-daily time scales. Our primary focus is here on the differences between these two temperature levels in the

context of extreme precipitation, winds, and storminess. The availability of high-frequency model output (3 hourly) allows us to investigate changes in sub-daily events and also to extract storm tracks using a tracking algorithm (Feser et al., 2015b).

5 The structure of the study is as follows: Section 2 describes the data and explains the methods used in the analysis. The impact of the horizontal resolution on the representation of atmospheric large-scale circulation is investigated in Section 3.1. The historical runs are validated against observed mean atmospheric circulation and precipitation, as well as high percentiles of daily precipitation in Section 3.2. Section 4 focuses on changes in the mean climate and weather extremes. A summary and discussion follows in Section 5.

## 10 2. Data and Methods

### 2.1 Data

To assess the importance of the horizontal model resolution, we first analyzed historical runs of CAM5.1 (<http://www.cesm.ucar.edu/models/cesm1.0/cam/>), provided by C20C+ Detection and Attribution Project (<http://portal.neresc.gov/c20c/main.html/>). We compare three runs, which cover the period 1979-2005 and  
15 are performed at different resolutions. The CAM5-1-2degree run (hereafter CAM5\_2, Wolski et al. 2014), the CAM5-1-1degree run (hereafter CAM5\_1, Stone et al. 2017) and the CAM5-1-0.25degree run (hereafter CAM5\_0.25, Wehner et al. 2015) are performed at atmospheric horizontal grid distances of  $2.5^{\circ} \times 1.875^{\circ}$ ,  $1.25^{\circ} \times 0.937^{\circ}$ , and  $0.3125^{\circ} \times 0.234^{\circ}$ , respectively. The 1979-2005 runs use historical values for all forcings (GHGs, ozone, volcanic aerosol, solar), **except land-use changes (set at year-1850) and**  
20 **without changes in non-volcanic aerosols, which adopt a year-2000 era repeated annual cycle.**

Projected climate change impacts on the mean climate state and on extreme weather are investigated based on model simulations with CAM5.1.2 (hereafter CAM5.1.2\_0.25) at the highest available  $\sim 0.25^{\circ}$  horizontal resolution. The simulations are part of the Half a Degree Additional warming, Prognosis, and Projected Impacts (HAPPI) experiment (Mitchell et al. 2017). The project is designed to provide model output data  
25 describing climate and weather changes under  $1.5^{\circ}\text{C}$  and  $2^{\circ}\text{C}$  levels of global warming, as compared to pre-industrial conditions (1861-1880). The design of HAPPI (Mitchell et al. 2017) provides three time slice experiments, using atmosphere-only models, to create large ensembles of 10-year simulations for the present climate (2006-2015) and potential future climate under  $1.5^{\circ}\text{C}$  and  $2^{\circ}\text{C}$  levels of warming (2106-2115). The two future run ensembles will hereafter be referred to the  $+1.5^{\circ}\text{C}$  and  $+2^{\circ}\text{C}$ , respectively.  
30 Observed forcing conditions include Sea Surface Temperatures (SSTs) and sea-ice (Taylor et al., 2012). SSTs in future scenarios are prescribed by summation of the observed 2006-2015 SSTs and an offset estimated between decadal-averages of the 2006-2015 period and the projected warmer global conditions for the 2091-2100 period. The 2006-2015 runs use 2006-2015 values for all forcings (GHGs, non-volcanic aerosols, ozone, volcanic aerosol, solar), except land cover (set at 1850 year). The Pathway 2.6 (RCP2.6,  
35 year 2095) is used to provide the model boundary conditions, including atmospheric greenhouse gas

concentrations, aerosols, ozone, land use, land cover for the 1.5°C scenario. For the 2°C scenario these conditions are the same, except the CO<sub>2</sub>, which are set to a weighted combination of the RCP2.6 and RCP4.5 for.

5 It is important to underline that the design of HAPPI future simulations use the same aerosol forcing (RCP2.6, year 2095). This protocol differs essentially from the protocol of historical simulations, causing a non-negligible decrease in the aerosol forcing in both future scenarios. Wehner et al. 2017 (accepted in ESD) found for these scenarios a remarkable reduction in total aerosol optical thickness over the Northern Hemisphere, reaching up to 50% over the North-Atlantic and European region. Thus the interpretation of differences between future and present climate could be complicated by the combined  
10 effects of the reduced aerosols and increasing CO<sub>2</sub>.

The simulated features of large-scale circulation are compared with reanalysis of monthly pressure at mean sea-level (hereafter SLP), winds at 850hPa level, and DJF precipitation rates (hereafter PR) for the period 1979–2005. For SLP and wind we use ERA-Interim, provided by the European Centre  
15 for Medium-Range Weather Forecasts (<https://www.ecmwf.int/en/research/climate-reanalysis/era-interim>), at the spatial resolution of ~0.75° x 0.75°. We use also NCEP-DOE Reanalysis 2 (Kanamitsu et al. 2002, <https://www.esrl.noaa.gov/psd/data/gridded/data.ncep.reanalysis2.html>), at 2.5° x 2.5° resolution. Precipitation is provided by the University of Delaware (V4.01), [http://climate.geog.udel.edu/~climate/html\\_pages/README.ghcn\\_ts2.html](http://climate.geog.udel.edu/~climate/html_pages/README.ghcn_ts2.html). It is a global gridded land data  
20 set, with the 0.5°x 0.5° horizontal resolution. For comparison of the large-scale features, all variables were interpolated on a common 2.5°x2.5° horizontal grid. For our analysis of daily precipitation data, we use E-OBS (Haylock et al., 2008; <http://www.ecad.eu>), provided by the European Climate Assessment and Dataset. The dataset contains daily precipitation sums on a 0.25° regular latitude–longitude grid for the period 1950–2015.

## 25 2.2 Methods

Our analysis of projected climate change focuses on the North Atlantic and European sector (27°–75° N, 80° W–45°E). While most of the analysis focuses on the DJF season, the analysis of storm tracks is extended to the period of October to March (ONDJFM).

We use the long historical run of CAM5.1 at ~0.25° resolution (CAM5\_0.25), which includes the 1979–  
30 2005 period, and also a five-member ensemble for the period 2006–2015 (CAM5.1.2\_0.25), when referring to present climate. Five-member ensemble simulations for the 1.5°C and 2°C level of warming are referred to as future +1.5°C and +2°C runs, respectively. Statistical significance of differences in the mean DJF climate between future and present climate is tested with the Wilcoxon signed rank test at the 5 % significance level. It is a non-parametric test and hence it can be used without the assumption that the  
35 population follows Gaussian distribution.

The analysis of the simulated large-scale circulation will be based on monthly means of hydro-meteorological variables for the winter (December, January, February, hereafter DJF) season.

Ambient flow over the North Atlantic is described by the meridional SLP gradient between the SLP in the vicinity of Azores and SLP over Iceland. The metric is relevant to the North Atlantic Oscillation index, hence the location and size of the regions are chosen to match the location of the simulated maxima and minima of SLP, i.e. 30°-20° W, 30°-40° N and 25°-15°W, 60°-70°N, in the present climate and future projections. Spatial patterns of the mean SLP fields were compared using centered pattern correlation. The maximum of the zonal wind was estimated for the region 0°-30° W, 50°-65° N.

The extreme precipitation analysis is based on the 95<sup>th</sup> percentiles of 3-h and daily total precipitation ratio and return values (RV) for a return period T=10 yrs. The RV were estimated by fitting Generalized Extreme Values (GEV) distribution by the method of maximum log-likelihood estimation (MLE) (Coles, 2001; Smith, 2003; Wilks, 2006, Gilleland and Katz, 2014) to a block (seasonal) maximum in the 50-yr sample of concatenated member runs. **The design of the HAPPI simulations satisfies requirements of stationarity and independence necessary to fit with a stationary GEV model.**

Return values (RV) for a given return period (T) are defined as values expected to be exceeded once per T-years. RVs are estimated as the values **corresponding to [(1 - 1/T)th quantile] of a sample** fitted to the GEV model. For example the 90<sup>th</sup> quantile (10 % exceedance probability) is an RV for a T=10-year period. The analysis here focuses on 10-year periods of RVs, because estimations for longer periods (e.g. 50-year periods with an exceedance probability of 2 %) is more prone to sampling errors and biases due to large uncertainties on the tails given relatively short samples.

The goodness-of-fit to the GEV model is estimated with the Anderson-Darling (A-D) test. The test is a modified version of the Kolmogorov– Smirnov goodness-of-fit-test. The A-D test gives more weight to the **tail and therefore is more suitable for EV distributions analysis. Analysis shows that it validates most of the estimations of extreme precipitation for mid-latitude and high latitudes. However, approximations for the regions in the southern parts of Europe, where the mean precipitation is much lower, have shown larger uncertainty. Similar results were obtained in the analysis of the extreme precipitation, where the Kolmogorov-Smirnov test was applied (Barcikowska et al. 2017).**

### 2.3 Storm tracks

Changes in storminess were explored with two measures of daily values during the DJF season. The first one uses high percentiles (95<sup>th</sup>) of daily wind speed. The second is a transient poleward temperature flux at 700 hPa, computed with the daily meridional wind and temperature deviations from the wintertime average. Anomalies were filtered with a 2-10 days bandpass (Butterworth) filter and averaged over the DJF season.

Storm tracks were extracted using a tracking algorithm according to Feser and von Storch (2008). The automated tracking approach facilitates the analysis of spatiotemporal variability of cyclones, their lifetime and intensity (Ulbrich et al. 2009; Neu et al. 2013). The algorithm consists of two parts: detection and tracking. The first part searches for the local minimum SLP and maximum wind speed. Additionally, before tracking, a spatial digital band-pass filter (Feser and von Storch, 2005) was applied to the 3-hourly output of SLP fields to extract mesoscale features of variability.

A storm was identified when a lifetime wind speed maximum exceeded  $10 \text{ m s}^{-1}$ , and a pressure minimum dropped to 995 hPa and a filtered pressure anomaly of  $-5 \text{ hPa}$ . Only tracks lasting more than 96 hours were taken into account in order to extract relatively long-lived and intense storms. Cyclones forming at latitudes higher than  $60^\circ \text{ N}$  were excluded, to align with the purpose of the study which focuses on the European climate.

5 Seasonal fields of spatial density (SPD) of 3-hourly storm occurrences were accumulated within a  $4^\circ \times 4^\circ$  grid and weighted by the unit area. Spatial intensity fields were computed by aggregating within  $3^\circ \times 3^\circ$  grid boxes the number of 3-h storm occurrences with maximum intensity exceeding certain thresholds. The threshold for the accumulated wind fields is  $10 \text{ m s}^{-1}$  and  $0.25 \text{ mm hr}^{-1}$  for precipitation. Additionally, 10 maximum intensity values were chosen from each  $3^\circ \times 3^\circ$  grid falling within an area of  $9^\circ \times 9^\circ$  from the center of the storm. This approach facilitates the analysis of the storm's impact not only in the regions with local maximum but also for the exposed regions within larger distances from the center.

### 3. Simulated winter mean climate and weather extremes

To evaluate the performance of the CAM5 simulation, we compare time-average (1979-2005) SLP fields 15 from observations with three CAM5 historical simulations each run at different resolution, where all data sets are interpolated to the lowest data set resolution ( $2.5^\circ \times 2.5^\circ$  lat-lon grid). ERA-I ( $\sim 0.75^\circ$ ) and NCEP-CFSR ( $\sim 0.34^\circ$ ) observations are provided at higher resolutions than NCEP/DOE 2 ( $\sim 2.5^\circ$ ), hence they better serve the purpose. The SLP fields in ERA-I and NCEP-CFSR are almost identical, with small differences over Greenland (not shown). Hodges et al. 2011 found also that these 20 reanalysis agree, both in terms of numbers and locations of extratropical cyclones, much better than the older ones (JRA-25) for both hemispheres and that intensities are higher. A comparison of ERA-I with NCEP/DOE 2 shows most differences in the vicinity of Greenland. The latter shows slightly higher SLP values over land and lower SLP values south-east of Greenland. Nevertheless, the SLP patterns share very high correlation (uncentered), which is 0.98. A shown below, the differences 25 between ERA-I and CAM5 are of larger magnitude than the observational differences.

Figure 1 shows that all simulations exhibit realistic patterns of the meridional SLP gradient. However, the gradient between the Icelandic Low and Azores High, which characterizes the typical North Atlantic Oscillation (NAO) pattern, intensifies with increasing resolution. The magnitude and the pattern in 30 ERA-I correlate best with the one simulated at similar horizontal resolution (CAM5\_1) ( $r=0.96$ ). Correlations with the remaining two are slightly smaller, i.e. 0.95 for CAM5\_2 and 0.94 for CAM5\_0.25. The magnitude in CAM5\_0.25 is most intense, which agrees well with stronger westerlies from Greenland towards the British Isles, indicating a stronger mid-latitude jet stream. Secondly, both CAM5\_2 and CAM\_1 show a strong positive SLP bias in the subtropical part of Europe and North Africa, 35 and negative bias extending from Iceland towards southeastern Europe and the Caspian Sea, causing the mean ambient flow (Fig 1 contours of differences between CAM5 and reanalysis) over the eastern N-

Atlantic and most of Europe to be more zonally oriented when compared to the reanalysis.

**The deficiencies in the SLP fields** are also reflected in the anomalies of zonal wind (Fig 2) along the borders of the SLP circulation patterns. Both, CAM5\_2 and CAM5\_1, exhibit anomalously strong westerlies extending across Europe from the British Isles towards Turkey. This corresponds with the zonal bias in the ambient flow and pattern of storm tracks, found in the same regions in CMIP5 models (Zappa et al. 2012). This zonal bias in the ambient flow (Fig 1) is strongly reduced in the high-resolution run (CAM5\_0.25).

The results presented here indicate that using high-resolution CAM5 simulations in applications to the winter climate over the Euro-Atlantic regions adds considerably better performance than simply spatially more detailed information. At higher resolution, the large-scale atmospheric flow and associated midlatitude jet stream is better represented, both in terms of the pattern and the magnitude. This improvement will presumably lead to a more realistic representation of the midlatitude storm tracks and associated with them wind and precipitation over Europe.

Figure 3a shows that mean seasonal precipitation in CAM5\_0.25 indeed bears a very close resemblance, compared to observations. However the comparison indicates also a much higher magnitude of precipitation over regions with complex orography (up to 1 mm day<sup>-1</sup>) such as the Alps and the west coasts of Scandinavia and UK. Our comparison of observed (EOBS) and simulated daily precipitation at the same resolution (~0.25°) also demonstrates very high skill of CAM5\_0.25 in simulating precipitation extremes. Figure 3b compares 90<sup>th</sup> percentiles of daily precipitation extremes, indicating that CAM5 skillfully captures the structure and sharp gradients over orographic complex subdomains. Again, in some mountainous regions like the northwest coast of the Balkan Peninsula and south-west coast of Turkey, the simulated values are much higher than the observed ones.

At the same time, it is important to note that constructing a homogenous and high-resolution observational data sets is severely limited over these regions. Thus the differences between these datasets may originate **either from the model bias or from observational bias** (deficient quality or lack of the observations in these regions). As pointed out for Spain, the differences between different observational datasets may be higher than differences between model simulations and a certain observational dataset (Gómez-Navarro et al. 2012).

Overall, the comparison strongly suggests that high-resolution runs provide a more accurate representation of the winter climate and weather for the Euro-Atlantic sector, where storms play an important role. A correct representation of storm tracks, governed by the ambient flow, is crucial for capturing the wind and precipitation extremes in the European region, thus in the following section we will focus on the analysis of CAM5 simulations at 0.25° x 0.25° horizontal grid.

### **3.1 Impacts of climate warming at the +1.5°C and +2°C temperature levels: large-scale atmospheric circulation and precipitation changes**

In this section, we investigate climate and weather changes associated with the two global warming temperature levels 1.5°C and 2°C, specified at the Paris climate agreement, and the recently experienced

climate. Differences in the forcing between two future sets of HAPPI experiments is confined to different CO<sub>2</sub> forcing and also to the SST offset, which corresponds to the difference between the decadal average of SSTs in the present climate and in projections reaching 1.5°C or 2°C levels of warming. Each of the experiment includes also internal climate SST variations (e.g. ENSO), being in the same phase i.e. during decade 2006-2015. Therefore it is expected that the impacts of internal variations will be canceled out while discriminating between all three experiments. The interpretation between the future and present climate is more complex in HAPPI experiments, as the latter includes also impacts resulting from reduced aerosol forcing. Unfortunately, the protocol of the project lacks an additional experiment, which would allow isolating these impacts. Thus at the moment we have to accept the possibility of different factors dominating the changes derived between “future vs present climate” and changes derived between the future scenarios. In the following part we will investigate future changes in the mean winter climate including precipitation and atmospheric circulation over the North Atlantic and Europe.

Here we explore the future response of winter large-scale circulation to the specified levels of global warming. Figure 4a,b depicts differences between the large-scale circulation at the 2°C level of warming (CAM5.1.2\_0.25) and the present climate. To aid interpretation of these changes in the context of the mean ambient flow, Figure 4a shows also the present climate SLP pattern, featured with the maximum in the vicinity of Azores, and minimum over Iceland. The average SLP difference between these two regions is estimated as 29.6 hPa and the interseasonal standard deviation within the ensemble is estimated as 6.5 hPa. At the 2°C warming level the meridional SLP gradient intensifies to 31.3 hPa, which is reflected in the positive SLP anomalies in the subtropical regions and the negative anomalies in the upper latitudes. The depicted pattern resembles to some extent the fingerprint of the previously found global warming response, characterized by intensified and poleward shifted meridional circulation cells and corresponding intensification and shift of the westerlies between these cells (Lu et al. 2007; Yin 2005; Bengtsson et al. 2006; Wu et al. 2011). Anomalously intense westerlies (Figure 4b) extend eastward from north of the British Isles to the north coast of Scandinavia. This feature corresponds to an increase in precipitation in these regions. The maximum change is located at the northwest coasts of the British Isles (up to ~0.8 mm day<sup>-1</sup>) and Norway (~1. mm day<sup>-1</sup>), which are directly exposed to the influence of extratropical cyclones and the associated large quantities of moisture. Precipitation increases slightly over northwestern Europe (France and Germany, up to 0.3 mm day<sup>-1</sup>). The intensification of the subtropical high (Fig 4b) is accompanied by easterly anomalies at the southern (equatorward) flanks of the anomalous divergent flow, which reduces precipitation, with a maximum near the center of the anticyclonic anomaly. The anomalies extend eastward and cover most regions of the Iberian Peninsula, but mostly they do not exceed a reduction stronger than 0.1 mm day<sup>-1</sup>. At the same time, the derived pattern is not entirely clear as it depicts positive SLP anomalies over Greenland, and reductions south of Greenland, very weak zonal wind and precipitation anomalies in the subtropics. In the comparison between future scenarios, the pattern seems more robust and of larger magnitude.

The difference, estimated between the two warming levels (Fig 5a) clearly depicts that the additional half a degree warming added to the +1.5°C level yields not only a remarkable intensification of the SLP gradient, but also a strong poleward shift of the circulation cells, midlatitude westerlies, and precipitation anomalies. The estimated SLP differences show statistical significance at 5 % level, with most regions showing non-zero changes (Fig S2). Figure 5 shows that the maximum SLP anomaly is located over the northern part of Bay of Biscay, while reduced precipitation expands north-eastward, through the Biscay Bay, France, southern parts of British Isles, and the North Sea. Drying over the northwest coast of the Iberian Peninsula is even stronger, compared to the difference in reference to the present climate. Therefore, the zone of increased precipitation is more confined towards the north, covering northern parts of the British Isles, and the Norwegian coast. **The changes in the large-scale circulation are also manifested in the meridional SLP gradient, which increases from 29.1 hPa to 31.3 hPa. It is worth noting that this difference is even larger than the one derived between the +2°C warming and the present climate.**

At the same time, changes associated with warming at the +1.5°C level are quantitatively and qualitatively different from the shown above. In fact, the derived changes (**Fig S1a**) in large-scale circulation manifest an opposite tendency, when compared with the previous results. However the magnitude of these changes is very small. **The meridional SLP gradient decreases from the 29.6 hPa to 29.1 hPa. Consistent with it are (Fig S1a) pronounced positive SLP anomalies over Greenland and negative SLP anomalies south-eastward of British Isles, which contributes to the weakening of the meridional cells.** These results explain also why the +2°C minus present pattern is much weaker in comparison to the one derived between future scenarios. **The derived discrepancy points again to the combination of competing factors i.e. reduced aerosols and increasing CO<sub>2</sub> which renders the interpretation the comparison of future and present climate more complex. This interpretation requires a separate analysis an additional, properly designed experiment to isolate the effect of aerosol reductions.** Thus in the following sections, we will focus mostly on the difference between the +1.5C and +2°C scenario.

### 3.2 Changes in daily and sub-daily precipitation extremes

In this section we investigate changes in daily and 3-h precipitation extremes associated with an increase in global warming from 1.5°C to 2°C (CAM5.1.2\_0.25). Precipitation extremes are **defined here as 95<sup>th</sup>** percentiles and 10-year return values, derived by fitting a GEV distribution to the HAPPI model outputs. Figure 6a,b presents the future response derived for 95<sup>th</sup> percentiles of daily and 3-h precipitation, associated with an additional half a degree of warming. The response shows a bipolar pattern, with an increase over the North Atlantic over the northern part of the typical midlatitude storm track region, and a decrease southward, over the region of anticyclonic anomalies. Significantly increased precipitation anomalies extend north-eastward from Nova Scotia through the northwest British Isles towards the Norwegian Sea and northern Scandinavia. The maximum change is located along the northwest coasts of the British Isles and Scandinavia (up to ~0.2 and 0.24 mm h<sup>-1</sup> in 3-h precipitation, respectively), which corresponds well with the derived changes in mean precipitation.

Figure 6 also exhibits a significant (at 5 % significance level) reduction over the Iberian Peninsula, north-western Europe, and southern flanks of the British Isles. The most radical decrease in **sub-daily** precipitation extremes occurs along the northwest coast of the Iberian Peninsula (-0.25 mm h-1) and in the vicinity of Biscay Bay (-0.18 mm h-1). It is worth noting that changes in extremes of **sub-daily** precipitation are larger and more significant over larger areas. For example, the local minimum found in the **extremes of sub-daily precipitation northwestward from the Iberian Peninsula, is less recognizable in daily precipitation extremes**, which may indicate a smaller impact of storminess to the daily scales, as compared to 3-h data.

The future response in 10-yr return values for sub-daily precipitation (Fig 6c), **derived from GEV statistical models, is consistent with the pattern derived from 95<sup>th</sup> percentiles and indicates even larger changes**. For example, the increase over the northwest coasts of the British Isles and north-western Scandinavia reaches up to 0.3 mm h-1. The decrease in the west part of the continent, found in the analysis of the percentiles, covers a larger area and extends more towards the center of the continent. The magnitude of the precipitation and their changes in the off-coastal areas is often smaller. Nevertheless, the percent changes (in reference to the climatological values at the 1.5°C level) (Fig 7) indicate pronounced decreases (approaching a 15 %) in the interior of France, over the North Sea, southern Scandinavia, southeast Europe and up to a 25 % increase in the interior of northeastern Scandinavia.

### 3.3 Climatology and changes in sub-daily wind extremes and storminess

In this section we investigate future changes in storminess associated with an increase in global warming from 1.5°C to 2°C. Apart from the chosen forcing scenario, additional uncertainty in predictions of future climate may be also related to the general model performance, known bias in the historical period and the ability to simulate certain features of interest. As such, a validation of the model skill in simulating the long-term climate is not necessarily a guarantee for skillful future projections. However, it is a useful indicator for the model's fidelity to reasonably simulate features of interest. Hence, before analyzing projected changes for the future, we will start the analysis of storminess here by focusing first on the long-term mean, simulated with CAM5\_0.25 for the period 1979-2005.

Here we use three different measures of storminess: the 95<sup>th</sup> percentile 3-h wind speeds, band-passed filtered transient poleward temperature flux (VT), and density of storm tracks, which are explicitly extracted with a tracking algorithm. All of these measures have certain limitations in characterizing storminess. Measures of wind extremes and transient temperature fluxes will not distinguish the cause of the changes, e.g. changing frequency or intensity of storms. An application of the Lagrangian approach facilitates extraction of storm tracks and their properties. However potential deficiencies of models in representing realistically storm features (e.g. underestimated intensity) often limit the feasibility of tracking algorithms to construct a representative sample of storms. Thus the robustness of that approach can be limited due to the sampling bias. An interpretation using all three measures facilitates a more complete description of the present climate and future changes in storminess.

The analysis of the historical run for the period 1979-2005 shows that CAM5\_0.25 reproduces the spatial

features of storminess very realistically compared to the observational-based data sets. For example, a strong meridional tilt is skillfully captured in all three measures (Fig 8b, Fig 9a and Fig S3). For VT (Fig 8b), not only the spatial pattern but also the intensity agrees remarkably well upon direct comparison with the observational climatology (<http://www.met.reading.ac.uk/~swrshaff/sstanom.html>). The VT pattern manifests the full spatial spectrum of the location of extratropical cyclone activity. The pattern spreads across the subtropical and midlatitude North Atlantic, featuring maximum values along the region from Newfoundland, across the east Atlantic between the British Isles and Iceland, to the Norwegian Sea. The simulated maximum intensity of VT yields approximately the value of  $25 \text{ }^\circ\text{C m s}^{-1}$ , which is very close to the derived values from the ECMWF Reanalysis. The simulated intensity with CAM\_0.25 is much more realistic in comparison with one of the CMIP3 models (<http://www.met.reading.ac.uk/~swrshaff/sstanom.html>), with typically much lower horizontal resolution.

**In the latter, the strength of the storm intensity was found to be nearly half of the observed one.**

**For high wind speed percentiles (Fig S3)**, which have been widely used (e.g. Krueger et al. 2013) as a simple measure of storm activity, CAM5\_0.25 reproduces the pattern of local maximum very closely to the one found in VT. The simulated intensities also bear a close resemblance to the wind extremes (not shown) in reanalysis data *i.e.* ERA-Interim and CFSR. CFSR, which has the finest ( $\sim 0.25^\circ\text{-}0.5^\circ$ ) horizontal resolution, shows a much better agreement with the model. The ERA data set shows lower values than CFSR, especially over the vicinity of the local maximum. The apparent difference stems most likely from the underestimation of midlatitude extreme winds in ERA-Interim and ERA-40, which appears to be related to their relatively coarser spatial and temporal resolution (Chawla et al., 2013; Pielke, 2002; Stopa and Cheung, 2014; Sterl and Caires, 2005; Campos and Guedes Soares, 2017). As in the case of precipitation mentioned previously, this points again towards the finding, that differences between different observational data products may be as large or even larger than deviations of climate simulation relative to a certain reference dataset (Gómez-Navarro et al. 2012). A coarse model resolution is however not the only explanation for too low wind speeds for high wind percentiles. As e.g. shown by Rockel and Woth (2007), even regional climate models simulate too low wind speeds for high percentiles if no gustiness correction is applied to the model output.

The climatology of the spatial track density (Fig 9a), derived from the tracking algorithm, agrees reasonably well with the tracks gleaned from observations (Zappa et al. 2012, **Zappa et al. 2013**, Hodges et al., 2003). However, the pattern in CAM5\_0.25 exhibits a maximum shifted towards the Norwegian Sea and does not manifest a strong activity in the region south-east of Greenland. This feature is well captured in the CAM5\_0.25 wind speed percentiles and is most likely associated with the short-lived katabatic winds that descend from the Greenland ice sheet. These features are however not of interest for our analysis and the tracking algorithm used in this study is tailored to extract only the long-lived and most intense cyclones. Upon visual inspection it can also be suggested that the track density simulated in CAM5\_0.25 is improved, as compared to the low-resolution CMIP3 and CMIP5 models. The CMIP models have been shown to exhibit a very strong zonal bias with positive anomalies in central Europe and negative values over the

Norwegian Sea (Zappa et al. 2012). We note however that the verification whether increasing the resolution improves the simulated climatology of midlatitude storms demands further analysis. This would require a unified methodology, with the same tracking algorithm applied to all of the datasets. Overall, first results shown here indicate that the CAM5\_0.25 reproduces features of storminess considerably more realistically than coarse resolution simulations, both, in terms of spatial pattern and intensity. This increases our confidence in the skill in projections of future storminess projected with the CAM5\_025, and is the focus of the remainder of this section.

Figure 8a depicts differences in the response between 1.5°C and 2°C level of warming, derived for the 95<sup>th</sup> percentile of 3-h wind speed. The derived changes show a bipolar pattern, similar to the one found for extreme precipitation. The most radical decrease in wind speeds manifests at the poleward fringe of the subtropics (40° N), between the Iberian Peninsula and the Azores. This region overlaps well with the location of maximum easterly anomalies at the southern flanks of the winter anticyclonic anomaly, found in the analysis of changes in general atmospheric circulation (see Fig 5). Thus it is likely that the simulated reductions in extreme winds are to a large extent caused by the poleward shift of the large-scale circulation, the signature of which is the weakening of the westerlies at the poleward flanks of the subtropics.

The response to the additional half a degree warming is also expressed as a remarkable increase in extreme winds over the northern half of the typical storm track region, with the maximum located between Iceland and the British Isles and along the Scandinavian coast. This feature is highly consistent with the response pattern derived for VT (Fig 8b). Changes in VT indicate a pronounced intensification of storminess on the poleward flanks of their DJF climatology, again featuring a maximum between Iceland and the British Isles and an eastward extension along the Scandinavian coast. Small negative anomalies occur over the Norwegian Sea, north-east of Iceland. A similar response is found in the storm track density (Fig 9a), showing an increase over the eastern North Atlantic and negative anomalies north-east of Iceland. Positive anomalies found in all measures of storminess collocate well with the local maximum of the intensification of the mean DJF westerlies (Fig 5), which is consistent with the eddy-driven nature of the midlatitude jet stream.

The analysis of the intensity accumulated along the extracted tracks provides further insights. Figure 9b shows an increase in the number of days, which exceed certain thresholds of precipitation and wind (0.25 mm h<sup>-1</sup> and 10 m s<sup>-1</sup>, respectively). The derived pattern shows similar features to those in the track density except that the positive changes are extended north-east of the Norwegian Sea. An additional analysis (not shown), repeated for higher thresholds of wind and precipitation, confirms the previous results in that it also exhibits an increase along the Scandinavian coast, indicating that the pattern becomes more zonal for higher intensities.

Overall, the increase manifested in the track density fields over the eastern North Atlantic, between the British Isles and Iceland, is consistent with the anomalies in VT. This suggests that the change in storm activity in this region is influenced by the increased frequency of storms. The increase in VT and in the number of high intensity days (as diagnosed from wind and precipitation) becomes clearly visible also over

the Norwegian Sea, despite no tendencies in track density in this area. For the increased thresholds of the intensity, positive anomalies emerge also at the coastal regions of Scandinavia, which are also accompanied with insignificant or zero tendencies in track density. Therefore the response found in storm activity over the Norwegian Sea could be alternatively explained by an increase of the intensity of the storms, rather than frequency. This is however a subject for a separate and more thorough analysis. It is also important to note that the storm tracks analyzed here exhibit a very strong year-to-year variability. Thus the statistics derived here may suffer from large uncertainty, and should be repeated when a larger number of ensemble simulations becomes available, in order to facilitate a reduction in the sampling error.

#### 4. Summary and Discussion

10 In this study we assess near-term regional winter climate and weather changes over the North Atlantic Ocean and Europe associated with the 1.5°C and 2°C levels of global warming. The design of most state-of-the-art experiments, e.g., Coupled Model Inter-comparison Project (CMIP), are not well suited to address questions on climatic changes associated with the specific climate policy goals. This is due to the fact that CMIP experiments are set in the framework of responses to the particular concentration scenarios, rather than to the particular level of warming. Therefore, we use here a set of ensemble simulations provided by the HAPPI project. The design of that experiment reduces the impacts of different phases of climate variations and thus facilitates differentiation of the climate response between the two warming levels. The CAM5 simulations provide a set of future climate experiments, describing the global climate and weather at  $\sim 0.25^\circ$  horizontal resolution and at **sub-daily** time-scale (3-h). Hence these simulations create a unique opportunity to explore changes and physical linkages between them across spatial and temporal scales. Additionally, a set of CAM5 historical simulations, provided at different horizontal resolutions, facilitates an insightful analysis of the benefits of increasing horizontal resolution in regional climate applications.

25 In the first part of our manuscript, we focused on the assessment of the model's ability to realistically represent key features of winter climate and weather over the Euro-Atlantic sector. Our analysis of the runs, performed at horizontal resolutions ranging from  $\sim 2^\circ$  to  $0.25^\circ$ , has shown a substantial improvement in simulated large-scale circulation, specifically the meridional SLP gradient and midlatitude zonal winds. The zonal bias of the ambient flow over the North Atlantic and Europe, common for low resolution CMIP3 and CMIP5 (Zappa et al. 2012) models, has been very clearly reduced with the highest model resolution. To a large extent, the reduction of the zonal bias may result from a much better skill to capture ambient flow-orographic interactions in the model with finer horizontal resolution, suggesting an important upscaling-impact of regional scales in shaping the large-scale circulation.

35 In the second part of the manuscript, we investigated near-future changes, associated with global warming at the temperature levels specified by Paris agreement. The pattern of the future response, when 2°C warming is compared to the present climate, confirms typical fingerprints of climate response. These are characterized by a poleward shift and intensification of the meridional circulation cells, manifested here as

strengthening meridional SLP gradient, and poleward strengthening and eastward extension of midlatitudes (Lu et al., 2007; Yin, 2005; Bengtsson et al., 2006; Wu et al. 2011; Feser et al. 2015).

5 However, different to previous studies, our analysis didn't identify a local maximum of anticyclonic SLP anomalies over the central Mediterranean. This feature was found in many CMIP3 and CMIP5 simulations (Giorgi and Lionello, 2008; Giorgi and Coppola, 2007; AR5, IPCC 2007) and was often used as an explanation (Giorgi and Lionello, 2008) for reduced precipitation in most parts of this region. Instead, in our analysis, the center of the anticyclonic anomaly is shifted north-westward, which locates it over the North Atlantic, north-westward of the Iberian Peninsula. This feature corresponds well with the shift in drying anomalies, which extend from the eastern North Atlantic and covers only western parts of  
10 Mediterranean.

The reason for this difference may be associated again with a strong positive bias in SLP over the Mediterranean and associated zonal bias of ambient flow, persisting in most of CMIP3 and CMIP5 models. Thus, the maximum of the SLP field over the Mediterranean might be partly an expression of that bias. Increasing horizontal resolution to  $\sim 0.25^\circ$  reduces the SLP bias almost completely, as shown in our  
15 analysis, which might explain the difference in the response pattern. In contrast to this result, other simulations using a  $\sim 0.5^\circ$  horizontal model resolution (Barcikowska et al. 2017) indicated a strong anticyclonic intensification and drying over most of the Mediterranean, despite remarkable reduction of the bias. Therefore, the explanation of this difference in the projected pattern may have other/or additional causes and demands further exploration running different models at different resolutions.

20 Our analysis also provides additional insights into the evolution of the response, as a function of changing global temperature and suggests that the poleward shift and intensification of the meridional circulation cells and midlatitude westerlies occurs mostly during the additional half a degree of warming beyond the  $1.5^\circ\text{C}$  level. The difference in the response between  $2^\circ\text{C}$  and  $1.5^\circ\text{C}$  levels is shifted poleward, compared to the changes estimated between  $2^\circ\text{C}$  and present climate. The maximum anticyclonic SLP anomaly is  
25 located over the Bay of Biscay, which corresponds well with strong relative drying in this region. These drying anomalies extend also further north-east towards the North Sea, shifting the borderline between opposite sign tendencies northwards. Maximum precipitation anomalies occur in the northwest parts of British Isles, along north-west coast of Scandinavia and the Norwegian Sea.

**The evolution of the future response shows a much stronger and distinct pattern compared to the**  
30 **changes prior to the  $1.5^\circ\text{C}$  level of warming. This** amplification in the change may hence be a reflection of the asymmetry in forcing changes between present climate and for the  $1.5^\circ\text{C}$  and  $2.0^\circ\text{C}$  experiments. The changes associated with warming at the  $1.5^\circ\text{C}$  level stem from an interplay of a number of forcings, including strong aerosol reductions, while an additional half a degree warming is solely a consequence of  $\text{CO}_2$  increases and ocean warming.

35 The response found here of winter weather over the North Atlantic and Europe is largely consistent with the changes found for the mean climate state and large-scale circulation. An increase in warming from  $+1.5^\circ\text{C}$  to  $2^\circ\text{C}$  level suggests a poleward intensification of daily and **sub-daily** extreme wind and

precipitation. These tendencies show the most pronounced impact in the regions most exposed to the inflow of moisture from the North Atlantic, e.g., the British Isles and northwest Scandinavia, where the 95<sup>th</sup> percentiles of 3-h precipitation increase up to 0.2 mm h<sup>-1</sup> and 0.24 mm h<sup>-1</sup>, respectively. The response pattern derived from daily precipitation shows a very similar pattern to the one derived from 3-h data.

5 However, the latter exhibits larger magnitudes and encompasses larger areas with significant changes. Changes derived with GEV approximations, indicating even more radical shifts, show an increase in 10-yr return levels of up to 0.3 mm h<sup>-1</sup> in the coastal regions of British Isles and North-west Scandinavia. The magnitude of changes in precipitation is smaller in the inland areas. However, many regions like northeast Scandinavia, may still be strongly impacted by an increase of up to 20 %, when compared to the 1.5°C

10 level. Consistent with changes in the mean precipitation along the southern coast of Scandinavia, the east coast of the British Isles and North Sea indicate a slight decrease. These tendencies are more intense and expand towards western Europe, exhibiting an up to 15 % decrease over France and exceeding a 25 % decrease over the interior and eastern Iberian Peninsula.

Derived changes in extreme precipitation and wind correspond well with changes in storminess, measured

15 here with the transient poleward temperature flux (hereafter VT) and features of explicitly extracted storm tracks. The projected future response, derived from sub-daily VT and from spatial density of the extratropical storm tracks, indicates an increase of storm activity towards the northern side of the current storm track (between Iceland and the British Isles) but also a decrease north-east of Iceland.

The decrease in storminess at the northern flanks of the storm track, measured as the frequency of intense

20 storms, has been identified in the CMIP5 projections (Zappa 2013). Similar to our analysis, the future response according to CMIP5 models suggests a polar amplification of global warming, associated strongly with the Arctic sea-ice loss. This in turn reduces the lower atmosphere meridional temperature gradient and also baroclinicity, shown here by the decrease of zonal wind northeastward from Iceland, which is consistent with the reduced storminess in this region. At the same time, the minimum of warming SSTs

25 over the North Atlantic could lead to increased surface atmospheric baroclinicity (Brayshaw et al., 2009; Woollings et al., 2012) and thus enhance storminess over the eastern North Atlantic.

An increase of transient poleward temperature flux is found also over the Norwegian Sea, along the Scandinavian coast, which collocates well with the local maxima of increase in extreme precipitation and wind. The density of storm tracks doesn't indicate any spatially coherent tendencies in this region.

30 However, the positive tendencies in this region emerge when the extreme precipitation and wind events, associated with the extracted storm tracks are analyzed. In these regions we found an increase in frequency of 3hrly storm occurrences with exceptionally high intensities. The strength of this tendency increases with the intensity of the extreme event, which suggests the possibility of increased frequency of more intense storms. These results should however be confirmed by a more elaborate analysis, specifically targeting

35 changes in storms, and is the subject of further research.

**Acknowledgements:** The authors are grateful to Ángel Muñoz and Alex Petrescu for helpful discussions.

MZ was supported through the Cluster of Excellence “CliSAP” (EXC177), Universität Hamburg, funded through the German Science Foundation (DFG). DAS and MFW were supported by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, under contract number DE-AC02-05CH11231.

5

**"The authors declare that they have no conflict of interest."**

**Submission for a Special Issue in Earth System Dynamics: The Earth system at a global warming of 1.5°C and 2.0°C**

## 10 References

- UNFCCC, Adoption of the Paris Agreement, FCCC/CP/2015/L.9/Rev.1, [http://unfccc.int/documentation/documents/advanced\\_search/items/6911.php?preref=600008831](http://unfccc.int/documentation/documents/advanced_search/items/6911.php?preref=600008831), 2017.
- Baeriswyl, P. A. and Rebetez, M.: Regionalization of precipitation in Switzerland by means of principal component analysis, *Theor. Appl. Climatol.*, 58, 31–41, <https://doi.org/10.1007/bf00867430>, 1997.
- 15 Ban, N., Schmidli, J. and Schär, C.: Heavy precipitation in a changing climate: Does short-term summer precipitation increase faster? *Geophysical Research Letters* 42(4), 1 165–1 172, doi: 10.1002/2014GL062588, 2015.
- Barcikowska, M., Kapnick, S.B., and F. Feser,: Impact of large-scale circulation changes in the North Atlantic sector on the current and future Mediterranean winter hydroclimate. *Climate Dynamics*,**
- 20 **doi:10.1007/s00382-017-3735-5, 2017**
- Bengtsson L., Hodges K.I., Roeckner E.: Storm tracks and climate change. *J Clim* 19(15): 3518–3543, 2006.
- Brayshaw, D. J., T. Woollings, and M. Vellinga.: Tropical and extratropical responses of the North Atlantic atmospheric circulation to a sustained weakening of the MOC. *J. Clim.*, 22, 3146–3155, 2009.
- 25 Coles, S. (2001), *An Introduction to Statistical Modeling of Extreme Values*, Springer, London.
- Collins, M., and Coauthors: Long-term Climate Change: Projections, Commitments and Irreversibility. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, T.F. Stocker et al., Eds., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA., 2013.
- 30 Campos, M. and Guedes Soares, C.: Assessment of three wind reanalyses in the North Atlantic Ocean, *Journal of Operational Oceanography*, 10:1, 30-44, DOI: 10.1080/1755876X.2016.1253328, 2017.
- Chawla, A., Spindler, D.M., Tolman H.L.: Validation of a thirty year wave hindcast using the Climate Forecast System Reanalysis winds. *Ocean Modell.* 70, 189–206, 2013.
- Della-Marta, P. M., and Pinto, J. G.: Statistical uncertainty of changes in winter storms over the North
- 35 Atlantic and Europe in an ensemble of transient climate simulations. *Geophys. Res. Lett.*, 36, L14703–L14703, doi:10.1029/2009GL038557, 2009.

- Donat, M. G., and Coauthors: Updated analyses of temperature and precipitation extreme indices since the beginning of the twentieth century: The HadEX2 dataset. *J. Geophys. Res. Atmos.*, 118, 2098–2118, doi:10.1002/jgrd.50150, 2013.
- D.J. Dokken; K.L. Ebi; M.D. Mastrandrea; K.J. Mach; G.-K. Plattner; S.K. Allen; M. Tignor and Midgley, P. (Eds.): *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation*. Chapter 3: Changes in climate extremes and their impacts on the natural physical environment. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 109-230, 2012.
- Drijfhout, S., and Coauthors.: Catalogue of abrupt shifts in Intergovernmental Panel on Climate Change climate models. *Proc. Natl. Acad. Sci. USA*, 112, E5777–E5786, doi:<https://doi.org/10.1073/pnas.1511451112>, 2015.
- Feser, F., Barcikowska, M., Krueger, O., Schenk, F., Weisse, R., and Xia, L.: Storminess over the North Atlantic and Northwestern Europe: A review. *Quarterly Journal of the Royal Meteorological Society*, 141, 350-382. doi:10.1002/qj.2364, 2015a.
- Feser, F., Barcikowska, M., Haeseler, S., Levebvre, C., Schubert-Frisius, M., Stendel, M., von Storch, H., and Zahn, M.: Hurricane Gonzalo and its Extratropical Transition to a Strong European Storm. *Bulletin of the American Meteorological Society*, 96(2012), 51-55. doi:10.1175/BAMS-D-15-00122.1, 2015b.
- Feser, F. and von Storch, H.: Regional modelling of the western Pacific typhoon season 2004. *Meteorol Z* 17(4):519–528, 2008.
- Feser, F. and H. von Storch: A Spatial Two-Dimensional Discrete Filter for Limited-Area-Model Evaluation Purposes. *Mon. Wea. Rev.*, 133, 1774–1786, <https://doi.org/10.1175/MWR2939.1>, 2005.**
- Fleig, A. K., Tallaksen, L. M., James, P., Hisdal, H. and Stahl, K., 'Attribution of European precipitation and temperature trends to changes in synoptic circulation', *Hydrology and Earth System Sciences* 19(7), 3 093–3 107, 2015.
- Geng, Q., and Sugi, M.: Possible Change of Extratropical Cyclone Activity due to Enhanced Greenhouse Gases and Sulfate Aerosols—Study with a High-Resolution AGCM. *J. Climate*, 16, 2262–2274, 2003.
- Gilleland, E. and R.W. Katz.: extRemes 2.0: An Extreme Value Analysis Package in R. *Journal of Statistical Software*. <https://www.rap.ucar.edu/staff/ericg/extRemes/extRemes2.pdf>, 2014.
- Giorgi, F.: Climate change hot-spots. *Geophys Res Lett* 33:L08707, doi:10.1029/2006GL025734, 2006.
- Giorgi, F. and Coppola, E.: European climate-change oscillation (ECO). *Geophys Res Lett* 34(21): L21703. doi:10.1029/2007GL031223, 2007.
- Giorgi, F. and Lionello, P.: Climate change projections for the Mediterranean region. *Glob Planet Chang* 63(2–3): 90–104, 2008.
- Giorgi, F., Coppola, E. and Raffaele, F.: A consistent picture of the hydroclimatic response to global warming from multiple indices: Models and observations', *Journal of Geophysical Research: Atmospheres* 119(20), 11 695–11 708 (doi: 10.1002/2014JD022238), 2014.
- Gómez-Navarro, J. J., Montávez, J. P., Jerez, S., Jiménez-Guerrero, P. and Zorita, E.: What is the role of the observational dataset in the evaluation and scoring of climate models? *Geophysical Research Letters*,

- 2012, 39, L24701, doi:10.1029/2012GL054206, 2012.
- Hodges, K.I., B.J. Hoskins, J. Boyle, and C. Thorncroft: A Comparison of Recent Reanalysis Datasets Using Objective Feature Tracking: Storm Tracks and Tropical Easterly Waves. *Mon. Wea. Rev.*, 131, 2012–2037, [https://doi.org/10.1175/1520-0493\(2003\)131<2012:ACORRD>2.0.CO;2](https://doi.org/10.1175/1520-0493(2003)131<2012:ACORRD>2.0.CO;2), 2003.
- 5 Hodges, K.I., R.W. Lee, and L. Bengtsson: A Comparison of Extratropical Cyclones in Recent Reanalyses ERA-Interim, NASA MERRA, NCEP CFSR, and JRA-25. *J. Climate*, 24, 4888–4906, <https://doi.org/10.1175/2011JCLI4097.1>, 2011
- Hov, Ø., Cubasch, U., Fischer, E., Höppe, P., Iversen, T., Kvamstø, N. G., Kundzewicz, Z. W., Rezacova, D., Rios, D., Duarte Santos, F., Schädler, B., Veisz, O., Zerefos, C., Benestad, R., Murlis, J., Donat, M.,
- 10 Leckebusch, G. C. and Ulbrich, U.: Extreme weather events in Europe: preparing for climate change adaptation, Norwegian Meteorological Institute, Oslo, 2013.
- IPCC, Climate change, the physical science basis. In: Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL (eds) Contribution of working group I to the fourth assessment report of the intergovernmental panel on climate change. Cambridge University Press, Cambridge, UK, 2007.
- 15 IPCC, Climate change 2014: synthesis report. In: Pachauri RK, Meyer LA (eds) Contribution of working groups I, II and III to the fifth assessment report of the intergovernmental panel on climate change. IPCC, Geneva, Switzerland, 151 pp, 2014.
- Kanamitsu, M., W. Ebisuzaki, J. Woollen, S. Yang, J.J. Hnilo, M. Fiorino, and G.L. Potter: **NCEP–DOE AMIP-II Reanalysis (R-2)**. *Bull. Amer. Meteor. Soc.*, 83, 1631–1644, [https://doi.org/10.1175/BAMS-83-11-](https://doi.org/10.1175/BAMS-83-11-1631)
- 20 **1631**, 2002
- Kendon, E. J., Roberts, N. M., Fowler, H. J., Roberts, M. J., Chan, S. C. and Senior, C. A.: Heavier summer downpours with climate change revealed by weather forecast resolution model' *Nature Climate Change* 4(7), 570–576 (doi: 10.1038/nclimate2258), 2014.
- Kitoh, A. and Endo, H.: Changes in precipitation extremes projected by a 20-km mesh global atmospheric
- 25 model. *Weather Clim Extremes* 11:41–52 (ISSN 2212-0947), 2016.
- Knutti, R., Rogelj, J., Sedláček, J. and Fischer, E. M.: A scientific critique of the two-degree climate change target. *Nat. Geosci.* 9, 13–18, 2016.
- Krueger, O., Schenk, F., Feser, F. and Weisse, R.: Inconsistencies between Long-Term Trends in Storminess Derived from the 20CR Reanalysis and Observations. *J. Climate*, 26, 868–874, <https://doi.org/10.1175/JCLI-D-12-00309.1>, 2013.
- Lambert, S. J.: The effect of enhanced greenhouse warming on winter cyclone frequencies and strengths. *J. Climate*, 8, 1447–1452, doi:10.1175/1520-0442(1995)008<1447:TEOEGW>2.0.CO;2, 1995.
- Lambert, S. J.: Changes in winter cyclone frequencies and strengths in transient enhanced greenhouse warming simulations using two coupled climate models. *Atmosphere-Ocean*, 42, 173–181, doi:
- 35 <http://dx.doi.org/10.3137/ao.420302>, 2004.
- Lambert, S. J. and Fyfe, J. C.: Changes in winter cyclone frequencies and strengths simulated in enhanced greenhouse warming experiments: results from the models participating in the IPCC diagnostic exercise.

- Clim. Dynam., 26, 713–728, doi:<http://dx.doi.org/DOI 10.1007/s00382-006-0110-3>, 2006.
- Leckebusch, G. C., Koffi, B., Ulbrich, U., Pinto, J. G., Spanghel, T. and Zacharias, S.: Analysis of frequency and intensity of European winter storm events from a multimodel perspective, at synoptic and regional scales. *Clim. Res.*, 31, 59–74, 2006.
- 5 Lu, J., Vecchi G.A., Reichler, T.: Expansion of the Hadley cell under global warming. *Geophys Res Lett* 34(6):L06805. doi:10.1029/2006GL028443, 2007.
- Maraun, D.: When will trends in European mean and heavy daily precipitation emerge?, *Environmental Research Letters* 8(1), 14004, doi: 10.1088/1748-9326/8/1/014004, 2013.
- Pielke R.A.: *Mesoscale meteorological modeling*. New York: Academic Press. 676 p, 2002.
- 10 Pinto, J. G., Spanghel, T., Ulbrich, U., and Speth, P.: Assessment of winter cyclone activity in a transient ECHAM4-OPYC3 GHG experiment. *Meteorol. Z.*, 15, 279–291, doi:10.1127/0941-2948/2006/0128, 2006.
- Pinto, J. G., Froehlich, E., Leckebusch, G. and Ulbrich, U.: Changing European storm loss potentials under modified climate conditions according to ensemble simulations of the ECHAM5/MPI-OM1 GCM. *Nat. Hazards Earth Syst. Sci.*, 7, 165–175, 2007.
- 15 Pinto, J.G., Spanghel, T., Fink, A., Leckebusch, G.C. and Ulbrich, U.: Factors contributing to the development of extreme North Atlantic cyclones and their relationship with the NAO. *Climate Dyn.*, 32, 711–737, 2009.
- Pinto, J. G., S. Zacharias, A. H. Fink, G. C. Leckebusch, and U. Ulbrich: Factors contributing to the development of extreme North Atlantic cyclones and their relationship with the NAO. *Clim. Dynam.*, 32, 711–737, doi:<http://dx.doi.org/DOI: 10.1007/s00382-008-0396-4>, 2009b:
- 20 **Priestley, M. D. K., Pinto, J. G., Dacre, H. F. and Shaffrey, L. C.: The role of cyclone clustering during the stormy winter of 2013/2014. *Weather*, 72: 187–192, 2017.**
- Duarte Santos, D., Schädler, F., Veisz, B., Zerefos, O., Benestad, C., Murlis, R., Donat, J. M., Leckebusch, G. C. and Ulbrich, U.: Extreme weather events in Europe: preparing for climate change adaptation, 25 Norwegian Meteorological Institute, Oslo, 2013.
- Schaller, N. et al.: Human influence on climate in the 2014 southern England winter floods and their impacts. *Nature Climate Change*, 6, 627–634, doi:10.1038/nclimate2927, 2016.
- Shaw, T.A., Baldwin, M., Barnes, E.A., Caballero, R., Garfinkel, C.I., Hwang, Y.-T., Li, C., O’Gorman, P. A., Rivière, G., Simpson, I. R., Voigt, A.: Storm track processes and the opposing influences of climate 30 change. *Nat. Geosci.* 9, 656–664, 2016.
- Schenk, F. and Stendel, M.: On the Problem of Homogeneity in Observations and Reanalysis Data. In: Quante, M. & Colijn, F. (Eds.): *North Sea Region Climate Change Assessment*, Springer International Publishing, 2016.
- Schiemann, R., M. Demory, L.C. Shaffrey, J. Strachan, P.L. Vidale, M.S. Mizieliński, M.J. Roberts, M.
- 35 Matsueda, M.F. Wehner, and T. Jung.: The Resolution Sensitivity of Northern Hemisphere Blocking in Four 25-km Atmospheric Global Circulation Models. *J. Climate*, 30, 337–358, 2017.
- Stahl, K., Hisdal, H., Hannaford, J., Tallaksen, L. M., van Lanen, H. A. J., Sauquet, E., Demuth, S.,

- Fendekova, M. and Jódar, J.,: Streamflow trends in Europe: Evidence from a dataset of near-natural catchments, *Hydrology and Earth System Sciences* 14(12), 2 367–2 382 (doi: 10.5194/hess-14-2367-2010), 2010
- 5 Stahl, K., Tallaksen, L. M., Hannaford, J. and van Lanen, H. A. J.: Filling the white space on maps of European runoff trends: Estimates from a multi-model ensemble, *Hydrology and Earth System Sciences* 16(7), 2 035–2 047 (doi: 10.5194/hess-16-2035-2012), 2012.
- Sterl, A., Caires, S.: *Climatology, variability and extrema of Ocean waves: the web-based KNMI/ERA-40 wave atlas*. *Int J Climatol*. 25:963–977, 2005.
- 10 Stone, D. A. and Risser, M. D. and Ang'elil, O. M. and Wehner, M. F. and Cholia, S. and Keen, N. and Krishnan, H. and O'Brien, T. A. and Collins, W. D.: A basis set for exploration of sensitivity to prescribed ocean conditions for estimating human contributions to extreme weather in CAM5.1-1degree, *Weather and Climate Extremes*, 2017, submitted.
- Stopa, J.E., Cheung K.F.: Intercomparison of wind and wave data from the ECMWF Reanalysis Interim and the NCEP Climate Forecast System Reanalysis. *Ocean Modell.* 75:65–83, 2014.
- 15 von Storch, H.; Feser, F.; Haeseler, S.; Lefebvre, C. & Stendel, M.: A violent midlatitude storm in Northern Germany and Denmark, 28 October 2013. - In: Herring, S. C.; Hoerling, M. P.; Peterson, T. C. & Stott, P. A. (Eds.): *Explaining Extreme Events of 2013 from a Climate Perspective*, Chapter 22. *Bulletin of the American Meteorological Society*, S76-S78, 2014.
- Ulbrich, U., Leckebusch, G.C. and Pinto, J.G.: *Theor Appl Climatol* 96: 117. 20 https://doi.org/10.1007/s00704-008-0083-8, 2009.
- Wang, X., Swail, V., Zwiers, F., Zhang, X. and Feng, Y.: Detection of external influence on trends of atmospheric storminess and northern oceans wave heights. *Climate Dynamics*, Springer Berlin/Heidelberg, 32, 189-203, 2009.
- 25 Wehner, M. F., K. A. Reed, B. Loring, D. Stone, and H. Krishnan. 2017. Changes in tropical cyclones under stabilized 1.5°C and 2.0°C global warming scenarios as simulated by the Community Atmospheric Model under the HAPPI protocols. *Earth System Dynamics*, submitted.
- Wehner, M., Prabhat, K. A. Reed, D. Stone, W. D. Collins, and J. Bacmeister: Resolution dependence of future tropical cyclone projections of CAM5.1 in the US CLIVAR Hurricane Working Group idealized configurations. *Journal of Climate*, 28, 3905-3925, 10.1175/JCLI-D-14-00311.1, 2015.
- 30 Wehner, M. F., Stone, D., Shiogama, H., Wolski, P., Ciavarella, A., Christidis, N., Krishnan, H.: Early 21st century anthropogenic changes in extremely hot days, as simulated by the C20C+ Detection and Attribution multimodel ensemble. Submitted to *Weather and Climate Extremes* (special C20C+ issue), 2017.
- Woollings, T.: Dynamical influences on European climate: An uncertain future. *Philos. Trans. Roy. Soc.*, 368A, 3733–3756, 2010.
- 35 Wolski, P. and Stone, D. and Tadross, M. and Wehner, M. and Hewitson, Attribution of floods in the Okavango Basin, Southern Africa, *J. Hydrol*, 511, 350--358, 2014.
- Woollings, T, Gregory, J.M., Pinto, J. G., Reyers, M. and Brayshaw, D.J.: Response of the North Atlantic

- storm track to climate change shaped by ocean–atmosphere coupling. *Nat. Geosci.*, 5, 313–317, 2012.
- Wu, Y., Ting M.F., Seager R., Huang H., Cane M: Changes in storm tracks and energy transports in a warmer climate simulated by the GFDL CM2.1 model. *Clim Dyn* 37(1–2):53–72. doi:10.1007/s00382-010-0776-4, 2010.
- 5 **Yang, X., G.A. Vecchi, R.G. Gudgel, T.L. Delworth, S. Zhang, A. Rosati, L. Jia, W.F. Stern, A.T. Wittenberg, S. Kapnick, R. Msadek, S.D. Underwood, F. Zeng, W. Anderson, and V. Balaji: *Seasonal Predictability of Extratropical Storm Tracks in GFDL’s High-Resolution Climate Prediction Model. J. Climate*, 28, 3592–3611, <https://doi.org/10.1175/JCLI-D-14-00517.1>, 2015**
- Yin, J.H.: A consistent poleward shift of the storm tracks in simulations of 21st century climate. **10 *Geophys Res Lett* 32(18): L18701. doi:10.1029/2005GL023684, 2005.**
- Yin, J., Schlesinger, M., and Stouffer, R.: Model projections of rapid sea level rise on the northeast coast of the United States. *Nature Geoscience*, 2(4), 262–266. doi:<https://doi.org/10.1038/ngeo462>, **2009.**
- Yin, J.H. A consistent poleward shift of the storm tracks in simulations of 21st century climate. **15 *Geophys Res Lett* 32(18):L18701. doi:10.1029/2005GL023684, 2005.**
- Zappa, G., Shaffrey, L.C., and Hodges, K.I.: The Ability of CMIP5 Models to Simulate North Atlantic Extratropical Cyclones. *J. Climate*, 26, 5379–5396, <https://doi.org/10.1175/JCLI-D-12-00501.1>, 2012.
- Zappa, G., Shaffrey, L.C., K.I. Hodges, P.G. Sansom, and D.B. Stephenson: A Multimodel Assessment of Future Projections of North Atlantic and European Extratropical Cyclones in the CMIP5 Climate Models. **20 *J. Climate*, 26, 5846–5862, <https://doi.org/10.1175/JCLI-D-12-00573.1>, 2013.**




Figure 1. Time-mean average of the DJF sea level pressure [hPa] over the period 1979-2005, regridded to  $2.5^\circ \times 2.5^\circ$  horizontal grid for ERA-Interim (ERA-I,  $\sim 0.75^\circ$  lat-lon original resolution), NCEP/DOE 2 ( $2.5^\circ$  lat-lon), and CAM5 at  $\sim 2^\circ$  (CAM5\_2deg),  $\sim 1^\circ$  (CAM5\_1deg),  $\sim 0.25^\circ$  (CAM5\_0.25deg) lat-lon resolution. Contours show a difference, in reference to a) NCEP-NCAR, b-d) ERA-Interim.




Figure 2. Time-mean average of the DJF zonal wind [ $\text{m s}^{-1}$ ] over the period 1979-2005, regridded to  $2.5^\circ \times 2.5^\circ$  horizontal grid for ERA-Interim (ERA-I,  $\sim 0.75^\circ$  lat-lon original resolution), and CAM5 at  $\sim 2^\circ$  (CAM5\_2deg),  $\sim 1^\circ$  (CAM5\_1deg),  $\sim 0.25^\circ$  (CAM5\_0.25deg) lat-lon resolution. Contours show the difference, in reference to ERA-Interim.




Figure 3. a) Time-mean average of DJF monthly means of precipitation ratio [mm day<sup>-1</sup>], averaged over period 1979-2005, in observations (Delaware, 0.5° resolution); and the CAM5\_0.25 model, smoothed to 0.5°x0.5° horizontal resolution; b) DJF daily precipitation 90<sup>th</sup> percentiles [mm day<sup>-1</sup>], averaged over period 1980-2005 in observations (EOBS, 0.25° resolution) and CAM5\_0.25 model.




Figure 4. Difference between +2°C (runs 2106-2115) and present climate ensemble means (2006-2015) in DJF a) sea level pressure [shaded, hPa] and wind vector at 850hPa [ $m s^{-1}$ ]. Contours show DJF sea level pressure in present climate ensemble, with a local maximum in the vicinity of Azores and minimum in the vicinity of Iceland; b) precipitation [ $mm day^{-1}$ ] and zonal wind [contours,  $m s^{-1}$ ] in CAM5\_0.25.






Figure 5. Difference between +2°C and 1.5°C ensembles in DJF a) sea level pressure [shaded, hPa] and wind vector at 850hPa [m s<sup>-1</sup>]. Contours show (as in Fig 4a) DJF sea level pressure in present climate, with a local maximum in the vicinity of Azores and minimum in the vicinity of Iceland; b) precipitation [mm day<sup>-1</sup>] and zonal wind [contours, m s<sup>-1</sup>] in CAM5\_0.25.



**Figure 6** Difference between +2°C and +1.5°C ensemble experiments for DJF a) 95<sup>th</sup> percentile of 3hrly precipitation, b) 95<sup>th</sup> percentile of daily precipitation [mm h-1], c) 10-yr return values in 3hr precipitation CAM5.1.2\_0.25. Percentiles and return values are derived from the samples with values 5 larger than 1 mm day-1. Regions in a) and b) are stippled, for differences significant at 10% level.



5 **Figure 7. Fractional change between +2°C and +1.5°C ensembles (ratio of the difference and the climatological mean in the +1.5°C experiment) for 10-yr return values of 3hrly precipitation [ $\times 100$  %] in CAM5.1.2\_0.25. Differences in precipitation were estimated for the values larger than 1 mm day<sup>-1</sup>.**




Figure 8. a) Difference between +2°C and +1.5°C ensemble experiments for DJF) 95<sup>th</sup> daily wind percentiles, b) DJF 700hPa transient poleward temperature flux in CAM5.1.2\_0.25. Contours show the climatology derived for period 1979-2005 (CAM5\_0.25) [ $^{\circ}\text{C m s}^{-1}$ ], Values over high orography are masked.

5




Figure 9. a) Difference (contours) between +2°C minus +1.5°C ensemble experiments, and mean climatology (shaded) in years 1979-2005, derived for number of 3-h storm occurrences accumulated within 4°x 4° grid boxes (number/decade); Climatology derived for period 1979-2005 is shaded. +2°C and +1.5°C ensembles constitute five decadal periods each, 1979-2005 period is missing one (1981) year.

b) Difference between: +2°C minus +1.5°C ensemble experiments, estimated for a number of 3-h occurrences with maximum larger than 0.25 [mm h<sup>-1</sup>], and wind larger than 10m s<sup>-1</sup>, (number/decade). Maximum values were chosen from 3-h precipitation data on 0.25°deg, which falls into 3°x 3° grid boxes. Differences were computed for grid boxes with number over the threshold at least 20/per decade in both experiments.

**Figure S1. Difference between a) +1.5°C; b) +2°C and present climate (2006-2015) for DJF sea level pressure [shaded, hPa] and wind vector at 850hPa [shaded, m s<sup>-1</sup>].**

**Figure S2. Difference between +2°C and +1.5°C experiments for DJF a) sea level pressure [shaded, hPa] in CAM5.1.2\_0.25 0.25°x0.25° lat-lon resolution. Regions are stippled, where the difference is significant at 5% level.**

**Figure S3. Winter (DJF) climatology of 95<sup>th</sup> percentile of daily winds, derived for period 1979-2005 in CAM5\_0.25.**

10

15

20

25