

## ***Interactive comment on “A bias-corrected CMIP5 dataset for Africa using CDF-t method. A contribution to agricultural impact studies” by Adjoua Moise Famien et al.***

**T. Iizumi (Referee)**

iizumit@affrc.go.jp

Received and published: 4 January 2018

General comments:

The presented study developed the bias-corrected CMIP5 GCM daily dataset using a combination of CDF-t method and WFDEI (and EWEMBI) and then compare with the ISI-MIP dataset which based on different bias-correction method and reference data (WFD). Some extreme climate indices (daily-based metrics) as well as maize yield simulated by a crop model were compared across different datasets to characterize the quality of the bias-corrected GCM daily dataset provided in this study. I respect the authors' efforts conducting this comprehensive analysis. Although this study analyzes

C1

only West Africa, their findings have implications especially for agricultural impact studies across the world. I only have a few concerns (listed below) and believe that most of them are minor. I recommend the acceptance of this manuscript with minor revision.

Specific comments:

1. Section 3.1. Although the detailed description of CDF-t method may be available in earlier study (Michelangeli et al. 2009, Vrac et al., 2012, 2016), a more completed explanation of key characteristics of the method in this section is unavoidable to make this manuscript stand-alone. Otherwise readers have to scratch around for. Particularly, it would be great if you could add an brief explanation whether the method forces the maximum (or minimum) value of a climatic variable in the future projection to be the same with that in historical period or not.

2. Section 3.3. I am not convinced whether the current design of the sensitivity analysis is appropriate to evaluate the sensitivity of correction to the length of calibration period. When the data in 1979-1996 are used as the calibration subset, those in 1997-2013 are used as the validation subset; this is fine. But, there is no independent validation subset when the data in 1979-2013 are used as the calibration subset. And it is easily expected that biases in bias-corrected data become the smallest when all available data are used as the calibration data. Therefore, the conclusion that the correction with the longest calibration period leads to the smallest biases is not examined using independent data. However, I think this part is not essential in this study. Removing or reanalysis are possible for this part.

3. Fig. 5. In GUICOAST JAS, ISI-MIP bias-corrected precipitation data distantly distributed from WFD. Why? This is unreasonable because most methods including the ISI-MIP bias-correction method forces GCM data in the historical period similar to the reference data (WFD). A plausible explanation is necessary.

4. Fig. 14. Two tendencies are observed in this figure. One is that the 95th percentile precipitation values from ISI-MIP data largely varied by GCM compared to the spread

C2

across the GCMs found in CDF-t method. Why? Another one is that in SAHEL JAS the IPSL-CM5A-LR data corrected by CDF-t method are relatively far from other GCMs of CDF-t data. A brief explanation is required.

5. P25L2. "It indicates also that WFD data and related bias-corrected simulations should not be used anymore." I think this is overstated. Please consider rephrasing or removal. I agree that a use of WFD leads to biased crop yields in crop model simulation mainly due to biases in solar radiation, as demonstrated in your analysis. However, simulated yield variability and/or projected future change in yields would not be affected in relative term when biases in solar radiation are a main issue (for instance, see lizumi et al., 2010). Reliable projection of yield change in absolute term is challenging, and therefore projected relative change in yield is still only a main source of information for adaptation planning and other application.

6. Section 4.3. The presentation of daily-based metrics is relatively not well organized in the current manuscript. The analysis and findings on the daily-based metrics themselves are useful, but not comprehensive compared to earlier study examining daily-based metrics (e.g., lizumi et al., 2017, JGR, doi:10.1002/2017JD026613). Why did you select a limited number of metrics for this analysis? More importantly, it seems that the importance of the analysis results is not equal to that of the analysis of crop model. The simulated maize yield is used as a metric in the current manuscript, as the daily-based metrics are did so; though the maize yield has much importance in the manuscript compared to the daily-based metrics. A justification to present a limited number of daily-based metrics is necessary.

#### Technical corrections

7. P1L2-3. Why has CDF-t method never been applied to Africa? Is that due to low availability of daily weather observations in that region? 8. P2L6. "robust" biases. Do you mean "persistent" biases? 9. P2L19. "high and robust biases". Do you mean "large and persistent biases"? 10. P3L24. "northern summer". Is this "northern hemispheric

C3

summer" or "boreal summer"? 11. P'L7. "anomalies". Probably, "differences" is more appropriate here. 12. Fig. 3. "observations". This should read "references". 13. P12L6. "standardized variance". This should read "standardized standard deviation" by definition of Taylor diagram. 14. P21L6. "GDHY" are a hybrid of FAO country yield data, satellite-derived crop-specific vegetation index and global crop datasets on crop calendar, harvested area and production shares achieved by different growing season. Subnational yield statistics are used to validate the grid-cell yield estimates, but not used as the input to estimate grid-cell yields. 15. P23L9. "Rsds" -> "rsds"? 16. Fig. 19. The label of y-axis should be replaced by "maize yield" instead of "crop yield" to be more precise. 17. Table 2. The caption of the table needs to include sufficient information to interpret the results presented in the table. Saying "see detail in the text" is not acceptable from the viewpoint of readability.

References lizumi, T., M. Nishimori, and M. Yokozawa, 2010: Diagnostics of Climate Model Biases in Summer Temperature and Warm-Season Insolation for the Simulation of Regional Paddy Rice Yield in Japan. *J. Appl. Meteor. Climatol.*, 49, 574–591, <https://doi.org/10.1175/2009JAMC2225.1>

Interactive comment on Earth Syst. Dynam. Discuss., <https://doi.org/10.5194/esd-2017-111>, 2017.

C4