

Interactive comment on “Irreversible ocean thermal expansion under negative CO₂ emissions” by Dana Ehlert and Kirsten Zickfeld

Anonymous Referee #2

Received and published: 11 June 2017

The author investigated the reversibility of ocean heat uptake and sea level due to thermal expansion by linearly increasing and then decreasing atmospheric CO₂ in the UVic ESCM. They found that the sea level continues for several decades after atmospheric CO₂ starts to decline and does not return to the pre-industrial level. They further tested the sensitivity of vertical sub-grid scale ocean mixing and concluded that, in most cases, both rise and decline rate of sea level increase with increasing vertical ocean mixing.

This paper is potentially interesting and focuses on an important topic in the context of the Paris Agreement. I recommend this manuscript to be published in Earth System Dynamics pending on minor revisions. I have detailed my comments below.

1. The authors may want to explore more about the path of ocean heat take and

storage beyond the AABW formation and AMOC. For example, either observations and coupled climate models show that anthropogenic warming enters the Southern Ocean along the downwelling branch of the residual MOC (Liu et al. 2016), and heat is mostly stored preferentially around 45S where surface waters are subducted to the north (Armour et al. 2016). The authors' results also reveal this path, and moreover, show that it is independent from the AABW formation. In Fig. 7, either the default mixing or other settings (e.g., B&L high, constant 1.0) show a warming tongue extending from surface around 45S to deep ocean. By contrast, cooling occurs along the shelf of the Antarctic as related to the AABW formation.

2. Page 12, lines 3-4 "AMOC responses to global warming have been well studied (Stocker and Schmittner, 1997; Rahmstorf, 2006; Meehl et al., 2007) and the decline in the AMOC under warming is due to surface warming and freshening of North Atlantic surface waters (Rahmstorf, 2006)." Herein the authors may want to notice that the AMOC response to global warming is also related to AMOC stability (Liu et al. 2017). The AMOC may moderately weaken or collapse in response to CO₂ increase, depending on which stability regime it resides in current climate.

3. Besides the global sea level change, I (also many other readers) am interested in the regional sea level change since it is critical to particular coastal regions. I am wondering whether the authors can make some discussions about this. One key is that a great change of ocean circulation, for example, the AMOC weakening or overshoot, can modulate the dynamic sea level in the Atlantic and Southern Ocean.

References

1. Liu, W., S.-P. Xie, Z. Liu and J. Zhu, 2017, Overlooked possibility of a collapsed Atlantic Meridional Overturning Circulation in warming climate. *Sci. Adv.*, 3, e1601666, doi:10.1126/sciadv.1601666.
2. Liu, W., S.-P. Xie and J. Lu, 2016, Tracking ocean heat uptake during the surface-warming hiatus. *Nature Commun.*, 7, 10926, doi: 10.1038/ncomms10926.

[Printer-friendly version](#)[Discussion paper](#)

3. Armour KC, J Marshall, J Scott, A Donohoe and ER Newsom, 2016, Southern Ocean warming delayed by circumpolar upwelling and equatorward transport. *Nature Geosci.*, 9, 549–554.

Interactive comment on *Earth Syst. Dynam. Discuss.*, <https://doi.org/10.5194/esd-2017-45>, 2017.

Printer-friendly version

Discussion paper

