

Terrassa, Barcelona, Spain, December 20, 2017

Manuscript number: ESD-2017-79

Title: Quantifying changes in surface air temperature dynamics over several decades

Authors: D. Zapalla, M. Barreiro and C. Masoller

We have answered the comments of the two referees:

https://editor.copernicus.org/index.php/esd-2017-79-AC1.pdf?mdl=msover_md&jrl=430&lcm=oc108lcm109w&acm=get_comm_file&ms=61479&c=131449&salt=6270021082060555853

https://editor.copernicus.org/index.php/esd-2017-79-AC2.pdf?mdl=msover_md&jrl=430&lcm=oc108lcm109w&acm=get_comm_file&ms=61479&c=131648&salt=6370064861278646899

Here we present the **List of changes** that we have done in the main manuscript, as well as the additional information that was included to the *Supporting Information*

- To take into account the comment of the first reviewer about the appropriateness of “comparing the data between two fixed time period,” in the Supporting information we include a new section where we compare relative variations computed in three different time intervals. We show that maps with very similar spatial structures are found, confirming the robustness of our findings.
- To take into account the comments of the two reviewers, we have re-written several paragraphs of the introduction.
- To take into account the comments of both reviewers, we have added to the significance analysis presented in the Supporting information, a percentile-based significance test.
- To take into account the comment of the first reviewer, we have modified the title and the text, removing the terms “interdecadal changes” and “large-scale patterns”.
- To take into account the comment of the first reviewer, we have revised the conclusions and re-worded the sentence “confirm this migration in the XX century”
- As suggested by the second reviewer, we include in the Supporting information a new section with an overview of the Hilbert Transform and two figures displaying simple examples. Here we also clarify the technical comments (detrending: we remove the linear trend, we explain how the unwrapping is done and explain how the 5% is chosen, and what we mean by “exactly reconstructs”).
- As suggested by the second reviewer, in the revised text we included a sentence about the quality of the reanalysis in the blue spot in the Arctic.
- We have re-written the sentence about "fast oscillations".

Quantifying changes in spatial patterns of surface air temperature dynamics over several decades

Dario A Zappalà¹, Marcelo Barreiro², and Cristina Masoller¹

¹Departament de Física, Universitat Politècnica de Catalunya, Rambla St. Nebridi 22, Terrassa 08222, Barcelona, Spain

²Departamento de Ciencias de la Atmósfera, Universidad de la República, Igua 4225, Montevideo 11400, Uruguay

Correspondence to: Cristina Masoller (cristina.masoller@upc.edu)

Abstract. We study daily surface air temperature (SAT) reanalysis in a grid over the Earth surface, to identify and quantify **changes in SAT dynamics** during the period 1979–2016. By analysing Hilbert amplitude and frequency we identify the regions where relative variations are most pronounced (larger than $\pm 50\%$ for the amplitude and $\pm 100\%$ for the frequency). Amplitude variations are interpreted as due to changes in precipitation or ice melting; frequency variations, to a northward shift of the inter-tropical convergence zone (ITCZ) and a widening of the rainfall band in the western Pacific Ocean. The ITCZ is the ascending branch of the Hadley cell and thus, by affecting the tropical atmospheric circulation, ITCZ migration has far reaching climatic consequences. As the methodology proposed here can be applied to many other geophysical time series, our work will stimulate new research that will advance the understanding of climate change impacts.

Copyright statement.

10 1 Introduction

The unprecedented intensification of weather extremes is motivating research aimed at understanding long-term climatic variations (Barreiro et al., 2008; Coumou and Rahmstorf, 2012; England et al., 2014; Cai et al., 2014; Turco et al., 2015) that can have profound socio-economic impacts (Ghil et al., 2011) and trigger complex ecological adaptation mechanisms (Lejeune et al., 2002; Beaumont et al., 2011; Gottfried et al., 2012; Bordeu et al., 2016).

15 **Quantifying variations in surface air temperature (SAT) dynamics over several decades is a challenging problem because of non-stationarity and the presence of trends, measurement noise, multiple time scales, memory and correlations in the data (Franzke, 2012; Massah and Kantz, 2016); in addition, reanalysis data can be unreliable (due to the lack of observational constrains in many geographical regions), and reanalysis time series are insufficiently long (as reanalysis starts at the beginning of the satellite era). These challenges have motivated the use, for climate data analysis, of data-driven approaches that have**
20 **been commonly used for investigating observed complex signals in other fields of science (e.g., neurological, physiological, financial, etc.). Univariate analysis tools that have been used to analyze SAT time series include detrended fluctuation analysis, fractional analysis, wavelet analysis, etc. Bivariate analysis and the complex network approach has also allowed to uncover**

inter-relations between SAT *anomalies* in different regions (Tsonis and Swanson, 2008; Donges et al., 2009; Barreiro et al., 2011). In this approach the *seasonal cycle* is removed to eliminate the influence of solar forcing and the links represent correlations (linear or nonlinear), or statistical similarities between SAT dynamics in different areas (Tirabassi and Masoller, 2016). On the other hand, changes in the SAT seasonal cycle have also been investigated, and a trend to reduced cycle amplitude has been detected in many regions (Stine et al., 2009; Qian et al., 2011; Dwyer et al., 2012; Stine and Huybers, 2012; Duan et al., 2017; Chambers et al., 2013; Wang and Dillon, 2014). However, changes in SAT dynamics over several decades (such as those observed in Fig. 1), have not yet been investigated at a global scale. In order to fill this gap, we use Hilbert analysis (described in the *Supporting Information*) to investigate SAT time series with daily resolution (reanalysis covering the Earth surface in the period 1979–2016). Our goal is to detect the most sensitive regions (“hotspots”) where variations in SAT dynamics over the last decades are more pronounced.

The Hilbert transform (HT) provides, for a real oscillatory time series, $x(t)$ with $t \in [1, T]$, an instantaneous amplitude, $a(t)$, and an instantaneous frequency, $\omega(t)$, for each data point of the time series, and thus allows to characterize how the amplitude and the frequency of a signal vary in time. If a signal does not have a sufficiently narrow frequency band, $a(t)$ and $\omega(t)$ will not have a clear physical meaning (Pikovsky et al., 2001). The usual solution is based on band-pass filtering to isolate a narrow frequency band; however, HT directly applied to the signal can still yield useful information. An alternative solution is based on the Hilbert-Huang Transform (Huang et al., 1998), that combines Hilbert analysis with the empirical mode decomposition, that decomposes an arbitrary real time series into components, each having the physical meaning of a rotation in the complex plane.

Because many natural geophysical time series have a seasonal periodicity, this has motivated the use of Hilbert analysis to characterize the time-varying oscillation amplitude, to investigate phase-shifts, phase-amplitude couplings, etc. Applications in various geophysical fields are discussed in Huang and Wu (2008). As more recent examples, Massei and Fournier (2012) used Hilbert analysis to characterize the daily variability of the Seine river flow from 1950 to 2008, uncovering linkages between river flow variability and global climate oscillations (the North-Atlantic Oscillation and the Madden-Julian Oscillation). Sun (2015) used Hilbert analysis to compute the daily phase shift between temperature signals recorded at the ground surface and at a depth of 5m in two meteorology stations in Taiwan from 1952 to 2008. Significant reductions in the phase shift from 1980s to 1990s were found, which was interpreted to be related to the warming of the Pacific Decadal Oscillation. Reddy and Adarsh (2016) applied Hilbert analysis to rainfall time series in India, and found that the multiscale components of rainfall series have similar periodic structure as global climate oscillations (the Quasi Biennial Oscillation, El Niño Southern Oscillation, etc.)

We have recently applied Hilbert transform to unfiltered daily SAT reanalysis (Zappalà et al., 2016). We have shown that the maps of time-averaged Hilbert frequency, $\langle \omega \rangle$, and of standard deviation, σ_ω , revealed well-defined large-scale structures which were consistent with known dynamical processes.

Here we use $a(t)$ and $\omega(t)$ to quantify SAT variations. Our hypothesis is that changes in $a(t)$ and $\omega(t)$, can yield information about variations in SAT dynamics. Specifically, we are interested in addressing the following questions: Which properties of $a(t)$ and $\omega(t)$ display relevant variations? Where are the regions where these variations are more pronounced? Which processes can be responsible of these variations? Can these variations be used as a quantitative measure of regional climate change?

5 2 Data

In the main text we present results from ERA-Interim daily SAT reanalysis (Dee et al., 2011), that covers the period from January 1979 to June 2016 with a spatial resolution of 2.5 degrees, both in latitude and in longitude. Thus, there are $N = 73 \times 144 = 10512$ geographical sites and in each site the SAT time series has $T = 13696$ days. In the *Supporting Information* we compare ERA-Interim with NCEP-DOE Reanalysis 2, that is an improved version of the NCEP Reanalysis I model (Kistler et al., 2001). It covers a longer time interval and has $94 \times 192 = 18048$ geographical sites. In order to perform a precise comparison between the results of the two datasets, in the NCEP-DOE Reanalysis 2 we consider the same time interval as the ERA-Interim dataset.

3 Methods

3.1 Hilbert analysis

15 To apply the Hilbert transform (described in the *Supporting Information*) we first pre-process each raw SAT time series, $r_j(t)$ (where $j \in [1, N]$ represents the geographical site and $t \in [1, T]$ represents the day): we eliminate the linear trend and normalize to zero mean and unit variance, obtaining $x_j(t)$. The Hilbert transform is then applied to $x_j(t)$, obtaining $y_j(t) = HT[x_j(t)]$. From $x_j(t)$ and $y_j(t)$, the amplitude $a_j(t)$ and the phase $\varphi_j(t)$ were calculated as: $a_j(t) = \sqrt{[x_j(t)]^2 + [y_j(t)]^2}$ and $\varphi_j(t) = \arctan[y_j(t)/x_j(t)]$. Taking into account the signs of $x_j(t)$ and $y_j(t)$, the phase is constrained to the interval $[-\pi, \pi)$. Whenever
20 an extreme value of the interval is reached, the phase jumps to the other end of the interval. By eliminating these sudden jumps (using a standard library function that appropriately adds $\pm 2\pi$) we “unwrapped” the phase obtaining a continuous variation in time, from which the frequency time series, $\omega_j(t)$, was obtained by calculating the derivative. Since the Hilbert algorithm (Bilato et al., 2014) gives errors in the extremes, in each time series we disregarded the initial and final 5% (see the *Supporting Information*). In this way, we have time series of length $T = 12328$ with the property that $a_j(t) \cos \varphi_j(t)$ exactly reconstructs
25 $x_j(t)$.

3.2 Measures used to quantify variations in SAT dynamics

Variations in the Hilbert amplitude were quantified by the relative change, $\Delta a / \langle a \rangle = (\langle a \rangle_1 - \langle a \rangle_f) / \langle a \rangle$, where $\langle a \rangle_f$ is the average value of the amplitude during the first 10 years of the time series (January 1979 to December 1988), and $\langle a \rangle_1$, during the last ten years (July 2007 to June 2016). Analogously, we calculated the relative change of amplitude variance, $\Delta \sigma_a^2 / \sigma_a^2$, of
30 average frequency, $\Delta \omega / \langle \omega \rangle$, and of frequency variance, $\Delta \sigma_\omega^2 / \sigma_\omega^2$. In the *Supporting information* we analyze how the spatial structures uncovered depend on the time-intervals used to calculate the relative variations: we compare with relative variations during the first and final five years of the reanalysis, and also, during the first half period and the second half period of the reanalysis. While the values of the relative variations vary with the time-interval considered, the spatial maps are remarkably robust as the same structures are found with the three time-intervals considered.

5 A similar analysis was performed to detect changes directly from the raw SAT time series, $r_j(t)$, by computing the amplitude of the *climatology* (or seasonal cycle), $c_j(t)$, and the variance of *anomaly* time series, $z_j(t)$.

Specifically, the amplitude of the climatology was calculated as: $a_j^{(\text{clim})}(I) = \max[c_j^I(t)] - \min[c_j^I(t)]$, where $c_j^I(t)$ is the climatology series calculated only in the time interval I . We remark that the climatology amplitude $a_j^{(\text{clim})}(I)$ is a scalar number that depends on the choice of the time interval I . We calculated the climatology amplitude in the first and last decade, as well
 10 as in the whole series. As before, we used these values to calculate the relative change $\Delta a^{(\text{clim})}/a^{(\text{clim})}$. Also, the variance of the anomaly time series $z_j(t)$ was calculated and then used to find the relative change, $\Delta\sigma_z^2/\sigma_z^2$.

With the goal of relating changes in Hilbert frequency with changes in statistical properties of SAT time series, an analysis of the number of zero-crossings was performed: for each $x_j(t)$ we counted the number of crossings through the mean value, $x = 0$. As with other quantities, we then calculated the relative change.

15 3.3 Significance Analysis

A statistical significance analysis was performed by surrogating Hilbert series. For each amplitude time series (i.e., in each grid point) 100 shuffle surrogates were generated and for each surrogate the relative change, $\Delta a^s / \langle a^s \rangle$, was calculated. Then, the average over the 100 surrogates, $\langle \Delta a^s / \langle a^s \rangle \rangle_s$, and its standard deviation, σ_s were used to define the significance threshold: the relative change computed from the original data was considered significant if it was higher than $\langle \Delta a^s / \langle a^s \rangle \rangle_s + 2\sigma_s$, or
 20 lower than $\langle \Delta a^s / \langle a^s \rangle \rangle_s - 2\sigma_s$. In the colour maps, regions where variations are not significant are displayed in white. The same test was applied to frequency variations and the other quantities, except for the climatology for which a surrogate test is not applicable. In the *Supporting Information* various thresholds are considered **and in addition, a non-parametric significance test is used. Here we present only the maps obtained with threshold $\pm 2\sigma$, because it is a compromise between uncovering the spatial regions where SAT changes are pronounced, and disregarding the areas where the variations are small.**

25 4 Results

We analyse the maps of $\langle \omega \rangle$, $\langle a \rangle$, σ_ω^2 and σ_a^2 , in the first ten years and in the last ten years of the period covered by the reanalysis, as well as the relative change between the two decades.

4.1 Analysis of Amplitude Variations

Figures 1(a) and 1(b) display $\langle a \rangle$ in the first and in the last ten years, respectively, and Fig. 1(c) displays the relative difference
 30 (see *Methods* for details). In Fig. 1(c) we see an area of large increase (more than 50%) of average amplitude, located in South America (red spot marked by a triangle), and an area of large decrease (again, more than 50%), located in the Arctic (blue spot marked by a circle). The raw SAT time series in these regions are displayed in Figure 2.

In both time series we clearly observe a change in the amplitude of the oscillations in the last ten years with respect to the first ten years, having a visual confirmation of the changes detected by Hilbert amplitude. The red spot in Amazonia, whose SAT series shown in Fig. 2(a) has an increasing amplitude, can be interpreted in terms of changes in precipitation. In particular,




Figure 1. Relative change of time-averaged Hilbert amplitude. (a) Amplitude averaged over the first ten years (January 1979 to December 1988). (b) Amplitude averaged over the last ten years (July 2007 to June 2016). (c) Relative change of Hilbert amplitude, $\Delta a / \langle a \rangle$. (d) Relative change of amplitude of the seasonal cycle, computed from the amplitude of the climatology, $\Delta a^{(clim)} / a^{(clim)}$. A good qualitative agreement is seen in the spatial structures in panels (c) and (d). Importantly, the structures uncovered by Hilbert amplitude are well defined, in comparison with those uncovered by the analysis of the climatology amplitude, which look noisier.

5 the increase of Hilbert amplitude is linked to the decrease of precipitation and to the lengthening of the dry season (as reported in (Gu et al., 2016; Liebmann et al., 2004; Fu et al., 2013)). This is due to the fact that, when precipitation decreases, the fraction of solar radiation that is not used for evaporation is used to heat the ground, which in turns heats the surface air. **This leads to higher extreme temperatures during the dry seasons, as can be observed in Fig. 2(a).** Regarding the blue spot in the Arctic region, where SAT series shown in Fig. 2(b) has a decreasing amplitude, it can be interpreted as due to the melting of sea

10 ice. In fact, when ice is present at the surface of the sea, it acts as an insulator, preventing heat exchange between sea and air. This causes a large amplitude of SAT cycle. On the other hand, if the ice melts, the air-sea heat exchange reduces the amplitude of the cycle. In particular, during winter the air temperature is mitigated by the sea and tends to have more moderated values. **It is important to take into account that this blue spot is in a region for which the observational constraints from satellites on the reanalysis are scarce, which decreases the quality of the reanalysis in the region. Therefore, in order to check whether the detected changes are robust, we performed the same analysis using the NCEP-DOE reanalysis dataset. The results, presented in the *Supporting Information*, confirm the presence of the blue spot in the Arctic.**




Figure 2. Surface air temperature in two regions where a clear change in the oscillation amplitude in the last ten years, with respect to the first ten years, is observed. (a) Site of coordinates (7.5 S, 307.5 E), marked with a triangle in Fig. 1(c). (b) Site of coordinates (75 N, 40 E), marked with a circle in Fig. 1(c).

5 Next, we compare the changes detected by Hilbert amplitude with those computed directly from SAT (by decomposing SAT time series into climatology and anomaly, as explained in *Methods*). Since the climatology term retains the seasonal variation, we expect its amplitude change to give similar indications as the Hilbert amplitude change. On the other hand, the anomaly term contains all the rapid variability, so we expect its variance to give similar results as the variance of Hilbert amplitude.

10 Figures 1(c) and 1(d), which display respectively the relative change of Hilbert amplitude and of climatology amplitude, and Figs. 3(a) and 3(b), which display respectively the relative change of Hilbert amplitude variance and of anomaly variance, confirm these expectations. The good qualitative agreement seen in the spatial structures in these maps confirms that Hilbert analysis directly applied to unfiltered SAT indeed gives a physically meaningful instantaneous amplitude, with average and variance values that are consistent with those computed from SAT.

5 In Figs. 3(a) and 3(b), however, there is a difference in the eastern Pacific Ocean, in the area marked with a circle. In particular, in Fig. 3(b) there is an area with large decrease of variance (deep blue, around -100%), while in (a) the decrease is less pronounced (light blue, around -65%) and extended over a smaller area. In addition, in Fig. 3(a) there is an orange-red area that indicates a moderate increase of variance (around 45%), while in (b) such area is absent. The reasons underlying these differences will be discussed later.




Figure 3. Relative change of amplitude fluctuations computed from the variance of (a) Hilbert amplitude, $\Delta\sigma_a^2/\sigma_a^2$; (b) the anomaly time series, $\Delta\sigma_z^2/\sigma_z^2$.

4.2 Analysis of Frequency Variations

Figure 4(a) displays the average frequency $\langle\omega\rangle$ in the first ten years, Fig. 4(b) in the last ten years, and Fig. 4(c) displays the relative change, $\Delta\omega/\langle\omega\rangle$. In Fig. 4(c) we note that in the eastern Pacific Ocean there are two small areas, enclosed by the circle, of intense increase (red) and decrease (blue) of frequency. They both represent frequency changes whose absolute values are larger than 100% and correspond to the same region where differences were detected in Fig. 3.

These two areas of opposite signs suggest that, between the initial and the final decade, there is a shift of the inter-tropical convergence zone (ITCZ) toward the north. The ITCZ involves strong convective activity, which causes rapid fluctuations of SAT, thus giving high values of instantaneous frequency, as shown in Figs. 4(a,b). Therefore, in the relative change of frequency, in regions corresponding to the initial position of the ITCZ we see a decrease, while in regions corresponding to the present position of the ITCZ we see an increase. For the same reason, the two red areas in the western Pacific Ocean (indicated by two squares) suggest an expansion of the tropical convective regions. **This interpretation is in agreement with previous works that have related a northward shift of ITCZ to an inter-hemispheric temperature gradient, as the one experienced during the last decades** (Yoshimori and Broccoli, 2008; Kang et al., 2009; Frierson and Hwang, 2012; Schneider et al., 2014; Talento and Barreiro, 2016). Regarding the red areas in the north Atlantic, in the north Pacific and in the south Pacific, they are consistent with an increase in the occurrence of fronts which cause large daily fluctuations of temperature and thus an increase of Hilbert frequency.

To gain insight into the physical meaning of the changes that are captured by Hilbert frequency, we use an alternative approach to estimate frequency variations: we define as “events” the zero crossings of SAT time series (Pikovsky et al., 2001) (detrended and normalised to zero-mean as described in *Methods*). Then, we count the number of events in the first ten years, in the last ten years, and calculate the relative variation.

Figure 4(d) displays the map of relative change of zero-crossings. We see that there is a qualitative good agreement with the spatial structures seen in Fig. 4(c), thus providing a physical interpretation for the observed variation of Hilbert frequency: the




Figure 4. Relative change of time-averaged Hilbert frequency (in units of oscillations/year). (a) Average in the first ten years (1979–1988). (b) Average in the last ten years (2007–2016). (c) Relative change of Hilbert frequency, $\Delta\omega/\langle\omega\rangle$. (d) Relative change of the number of zero-crossings of the normalised SAT time series. In (a) and (b) the colour scale is adjusted to represent in white the regions where the average frequency is one oscillation per year. In (c) and (d), a good qualitative agreement of spatial structures is seen; however, we note that Hilbert frequency detects stronger variations than those measured by the number of zero-crossings.

areas where the frequency increases (decreases) correspond to areas where the number of zero-crossings increases (decreases). We note that the relative variations in Hilbert frequency are more pronounced than those in the number of crossings, and this specifically holds in the regions where frequency variations are interpreted in terms of ITCZ migration.

10 Figures 5(a) and 5(b) display SAT time series in the dipole region indicated with the circle in Fig. 4(c), and also indicate (in red) the zero-crossings. We can understand the difference that was detected in this region between the variance of Hilbert amplitude (Fig. 3a) and the variance of anomaly (Fig. 3b). This difference is explained in the following terms: in the first decade the seasonal cycle is more irregular than in the last decade, probably a consequence of an El Niño event in 1982–1983. The anomaly series contains these slow fluctuations as well as the rapid ones, and thus its variance is affected by both effects. In contrast, the Hilbert amplitude is less affected by the slow fluctuations as its variance captures mainly the rapid fluctuations
5 of SAT.




Figure 5. Normalised SAT time series and number of zero-crossings in the regions indicated with a circle in Fig. 4(c). In the red region (2.5 N, 245 E), panel (a), the number of zero-crossings increases in the last ten years with respect to the first ten years (289/202 respectively), while in the blue region (7.5 S, 250 E), panel (b), it decreases (128/258 in the last/first ten years).

To demonstrate the robustness of our findings, in the *Supporting Information* we compare the results obtained from ERA-Interim with those obtained from another reanalysis dataset, NCEP-DOE. We find a good qualitative agreement in the spatial structures in the maps of $\langle \omega \rangle$, $\langle a \rangle$, σ_ω^2 and σ_a^2 , but we discuss also some relevant differences. In addition, to further understand the relationship between statistical properties of SAT and those of Hilbert amplitude and frequency, in the *Supporting Information* we apply Hilbert analysis to synthetic data generated by an autoregressive AR(1) process. We chose an AR(1) process because it is commonly used in the literature to model climate data. We find that, when increasing the noise intensity in the synthetic series, the Hilbert amplitude decreases while the frequency increases and show that this trend is also observed in real SAT time series.

5 Conclusions

We have used Hilbert analysis to quantify the changes in SAT dynamics, in a global scale, that have occurred over the last three decades. From the SAT time series with daily resolution we derived the amplitude and the frequency time series, and then calculated the relative change (between the first and the last decade) of average and variance of these series. Large variations of Hilbert amplitude (more than 50%) in the Arctic and in Amazonia were interpreted respectively as due to ice melting and

precipitation decrease. The analysis of Hilbert frequency also uncovered areas of large changes. In particular, two areas of opposite changes in eastern Pacific Ocean and two areas of increase in western Pacific Ocean suggest a shift towards north and a widening of the ITCZ. While there is evidence that ITCZ has moved north-south in the past, **to the best of our knowledge our work is the first one to confirm this migration in the last decades**. Our findings have important implications because, as the

10 ITCZ is the ascending branch of the Hadley cell, its migration affects both the Earth's radiative balance and the release of latent heat that drives the tropical atmospheric circulation. Taken together, these effects have not only local but far reaching climatic consequences. Additional analysis provided in the *Supporting Information* confirms the robustness of these observations.

As the methodology used here can be applied to many other climatological time series that exhibit well defined oscillatory behaviour, we believe that our work will stimulate new research to identify and quantify the impacts of climate change, directly

15 from observed data.

Code and data availability. The Hilbert algorithm used is available in Bilato et al. (2014); the data sets used are ERA-Interim Reanalysis, provided by the European Centre For Medium-Range Weather Forecasts (ECMWF), Reading, UK, from their website: <https://www.ecmwf.int> and NCEP-DOE Reanalysis 2, provided by NOAA Boulder, Colorado, USA, from their website: <http://www.esrl.noaa.gov/psd/>.

Competing interests. The authors declare no conflict of interest.

5 *Acknowledgements.* This work was supported in part by the LINC project (EU-FP7-289447), ICREA ACADEMIA (Generalitat de Catalunya, Spain), and by Spanish MINECO/FEDER (FIS2015-66503-C3-2-P). D. A. Zappalà also thanks European Social Fund and the Generalitat de Catalunya for the FI-AGAUR scholarship.

References

- Barreiro, M., et al.: Inferring long memory processes in the climate network via ordinal pattern analysis, *Chaos*, 21, 013101, 2011.
- 10 Barreiro, M. et al.: Abrupt climate changes: How freshening of the northern Atlantic affects the thermohaline and wind-driven oceanic circulations, *Annu. Rev. Earth Planet. Sci.*, 36, 33–58, 2008.
- Beaumont, L., et al.: Impacts of climate change on the world’s most exceptional ecoregions, *Proc. Natl. Acad. Sci.*, 108(6), 2306–2311, 2011.
- Bilato, R., O. Maj, and M. Brambilla: An algorithm for fast Hilbert transform of real functions, *Adv. Comput. Math.*, 40, 1159–1168, 2014.
- Bordeu, I., et al.: Self-replication of localized vegetation patches in scarce environments, *Sci. Rep.*, 6, 33703, 2016.
- 15 Cai, W., et al.: Increasing frequency of extreme El Niño events due to greenhouse warming, *Nat. Clim. Change*, 4(2), 111–116, 2014.
- Chambers, L., et al.: Phenological changes in the southern hemisphere, *PLoS ONE*, 8(10), e75514, 2013.
- Coumou, D., and S. Rahmstorf: A decade of weather extremes, *Nat. Clim. Change*, 2(7), 491–496, 2012.
- Dee, D. P. et al.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q.J.R. Meteorol. Soc.*, 137, 553–597, 2011.
- 20 Donges, J., et al.: The backbone of the climate network, *EPL*, 87(4), 48007, 2009.
- Duan, J., et al.: Weakening of annual temperature cycle over the tibetan plateau since the 1870s, *Nat. Comm.*, 8, 14008, 2017.
- Dwyer, J., M. Biasutti, and A. Sobel: Projected changes in the seasonal cycle of surface temperature, *J. of Climate*, 25, 6359–6374, 2012.
- England, M. et al.: Recent intensification of wind-driven circulation in the pacific and the ongoing warming hiatus, *Nat. Clim. Change*, 4(3), 222–227, 2014.
- 25 Franzke, C.: Nonlinear trends, long-range dependence, and climate noise properties of surface temperature, *J. of Clim.*, 25(12), 4172–4183, 2012.
- Frierson, D. M. W., and Y. T. Hwang: Extratropical influence on ITCZ shifts in slab ocean simulations of global warming, *J. of Clim.*, 25, 720–733, 2012.
- Fu, R., et al.: Increased dry-season length over southern amazonia in recent decades and its implication for future climate projection, *Proc. Natl. Acad. Sci.*, 110(45), 18110–18115, 2013.
- 30 Ghil, M., et al.: Extreme events: dynamics, statistics and prediction, *Nonlin. Processes Geophys.*, 18, 295–350, 2011.
- Gottfried, M., et al.: Continent-wide response of mountain vegetation to climate change, *Nat. Clim. Change*, 2(2), 111–115, 2012.
- Gu, G., et al.: Long-term changes/trends in surface temperature and precipitation during the satellite era (1979–2012), *Clim. Dyn.*, 46, 1091–1105, 2016.
- 35 Huang, N., et al.: The empirical mode decomposition and the Hilbert spectrum for nonlinear and non-stationary time series analysis, *Proc. R. Soc. Lond. A*, 454, 903–995, 1998.
- Huang, N. E., and Z. Wu: A review on Hilbert-Huang transform: Method and its applications to geophysical studies, *Rev. Geophys.*, 46, 8755–1209, 2008.
- Reddy, M.J., and S. Adarsh: Time-frequency characterization of sub-divisional scale seasonal rainfall in India using the Hilbert-Huang transform, *Stoch. Environ. Res. Risk Assess.*, 30, 1063–1085, 2016.
- Kang, S. M., et al.: The tropical response to extratropical thermal forcing in an idealized GCM: The importance of radiative feedbacks and convective parameterization, *J. Atmos. Sci.*, 66, 2812–2827, 2009.
- 5 Kistler, R., et al.: The NCEP-NCAR 50-year reanalysis: Monthly means CD-ROM and documentation, *B. Am. Meteorol. Soc.* 82, 247–268, 2001.

- Lejeune, O., et al.: Localized vegetation patches: A self-organized response to resource scarcity, *Phys. Rev. E*, 66(1), 010901, 2002.
- Liebmann, B., et al.: An observed trend in central south american precipitation, *J. of Clim.*, 17(22), 4357–4367, 2004.
- 10 Massah, M., and H. Kantz: Confidence intervals for time averages in the presence of long-range correlations, a case study on earth surface temperature anomalies, *Geophys. Res. Lett.*, 43(17), 9243–9249, 2016.
- Massei, N., and M. Fournier: Assessing the expression of large-scale climatic fluctuations in the hydrological variability of daily seine river flow (France) between 1950 and 2008 using Hilbert-Huang transform, *J. Hydrol.*, 448, 119–128, 2012.
- Pikovsky, A.; Rosenblum, M.; Kurths, J. *Synchronization: A universal concept in nonlinear sciences*. Cambridge University Press (2001).
- 15 Qian, C., et al.: Changes in the amplitude of the temperature annual cycle in china and their implication for climate change research, *J. of Climate*, 24(20), 5292–5302, 2011.
- Schneider, T., et al.: Migrations and dynamics of the intertropical convergence zone, *Nature*, 513, 45–53, 2014.
- Schwabeddar, J., and H. Kantz: Optimal extraction of collective oscillations from unreliable measurements, *Phys. Rev. Lett.*, 116, 104101, 2016.
- 20 Stine, A., and P. Huybers: Changes in the seasonal cycle of temperature and atmospheric circulation, *J. of Climate*, 25(21), 7362–7380, 2012.
- Stine, A., et al.: Changes in the phase of the annual cycle of surface temperature, *Nature*, 457(7228), 435–440, 2009.
- Sun, Y.-Y.: Instantaneous phase shift of annual subsurface temperature cycles derived by the Hilbert-Huang transform, *J. Geophys. Res. Atmos.*, 120, 1670–1677, 2015.
- Talento, S., and M. Barreiro: Simulated sensitivity of the tropical climate to extratropical thermal forcing: tropical SSTs and African land
- 25 surface, *Clim. Dyn.*, 47(3–4), 1091–1110, 2016.
- Tirabassi, G., and C. Masoller: Unravelling the community structure of the climate system by using lags and symbolic time-series analysis, *Sci. Rep.*, 6, 29804, 2016.
- Tsonis, A., and K. Swanson: Topology and predictability of El Niño and La Niña networks, *Phys. Rev. Lett.*, 100(22), 228502, 2008.
- 265 Turco, M., et al.: Observed climate change hotspots, *Geophys. Res. Lett.*, 42(9), 3521–3528, 2015.
- Wang, G., and M. Dillon: Recent geographic convergence in diurnal and annual temperature cycling flattens global thermal profiles, *Nat. Clim. Change*, 4, 988–992, 2014.
- Yoshimori, M., and A. J. Broccoli: Equilibrium response of an atmosphere-mixed layer ocean model to different radiative forcing agents: Global and zonal mean response, *J. of Clim.*, 21, 4399–4423, 2008.
- 270 Zappalà, D. A., et al.: Global atmospheric dynamics investigated by using Hilbert frequency analysis, *Entropy*, 18(11), 408, 2016.