



1 Modelling forest plantations for carbon uptake 2 with the LPJmL dynamic global vegetation 3 model

4 Maarten C. Braakhekke^{1,2,*}, Jonathan C. Doelman², Peter Baas³, Christoph Müller⁴, Sibyll Schaphoff⁴,
5 Elke Stehfest², Detlef P. van Vuuren²

6 ¹ Wageningen Environmental Research, P.O. Box 47, 6700 AA, Wageningen, the Netherlands

7 ² PBL Netherlands Environmental Assessment Agency, The Hague, The Netherlands

8 ³ Geoscience & Remote Sensing, Delft University of Technology, Delft, the Netherlands

9 ⁴ Potsdam Institute for Climate Impact Research, Potsdam, Germany

10 * Corresponding author: maarten.braakhekke@gmail.com

11

12 Abstract

13 We present an extension of the dynamic global vegetation model LPJmL to simulate planted forests
14 intended for C sequestration. We implemented three functional types to simulate plantation trees in
15 temperate, tropical, and boreal climates. The parameters of these functional types were optimized to
16 fit target growth curves (TGCs). These curves represent the evolution of stemwood C over time in
17 typical productive plantations and were derived by combining field observations and LPJmL estimates
18 for equivalent natural forests. While the calibrated model underestimates stemwood C growth rates
19 compared to the TGCs, it represents substantial improvement over using natural forests to represent
20 afforestation. Based on a simulation experiment in which we compared global natural forest versus
21 global forest plantation, we found that forest plantations allow for much larger C uptake rates on the
22 time scale of 100 years, with a maximum difference of a factor 1.9, around 54 years. In subsequent
23 simulations for an ambitious but realistic scenario in which 650 Mha (14% of global managed land,
24 4.5% of global land surface) is converted to forest over 85 years, we found that natural forests take up
25 37 PgC versus 48 PgC for forest plantations. Comparing these results to estimations of C sequestration
26 required to achieve the 2°C climate target, we conclude that afforestation can offer a substantial
27 contribution to climate mitigation. Full evaluation of afforestation as a climate change mitigation
28 strategy requires an integrated assessment which considers all relevant aspects, including costs,
29 biodiversity, and trade-offs with other land-use types. Our extended version of LPJmL can contribute
30 such an assessment by providing improved estimates of C uptake rates by forest plantations.



1 Introduction

It is increasingly clear that the stringent climate targets of the Paris Agreement cannot be achieved without negative emissions, i.e. net removal of carbon (C) from the atmosphere, later during the 21st century to compensate for emissions in the first half of the century (Gasser et al., 2015; Rogelj et al., 2018). Of the many proposed techniques to achieve C uptake, the two options currently most discussed for large-scale implementation are bioenergy in combination with carbon capture and storage and afforestation (Williamson, 2016). Both approaches will require considerable amounts of land and thus compete with other land-use functions, for example food production and biodiversity. While bioenergy is receiving considerable attention (van Vuuren et al., 2013), less consideration has been given to afforestation as a tool for land-based mitigation. C uptake occurs when natural vegetation is allowed to grow back on former croplands and pasture. While deliberately taking cropland or pasture out of production may involve costs, the direct management costs of natural regrowth are negligible. The carbon uptake rate of such natural regrowth, however, will usually achieve only a fraction of the potential C uptake rate at short time scales. Considerably higher C uptake rates are possible by planting forests (Paquette and Messier, 2010). Assisting regrowth by planting trees can substantially boost growth rates compared to natural forests because initial stages of primary succession (with herbaceous or shrub vegetation) are skipped and because fast-growing tree species can be selected. Moreover, trees are usually planted as saplings, cultivated under controlled conditions, which improves chance of successful establishment compared to development from seeds (Gladstone and Thomas Ledig, 1990).

Assessing the potential of land-based approaches for climate mitigation requires reliable estimates of C sequestration rates. Process-based models, such as dynamic global vegetation models (DGVMs), are a crucial tool for providing these estimates. DGVMs simulate carbon stocks and fluxes based on mechanistic descriptions of underlying processes, such as photosynthesis and organic matter decomposition in relation to environmental conditions. However, since the focus of DGVM development has traditionally been on natural ecosystems, very few of these models have an explicit representation of planted forests. Therefore, previous modelling studies on large-scale afforestation represented afforestation as natural regrowth (Krause et al., 2017), in some cases applying corrections to account for higher growth rates (Humpenoder et al., 2014; van Minnen et al., 2008).

In this paper we present an updated version of the DGVM LPJmL (Bondeau et al., 2007; Schaphoff et al., 2013), modified to explicitly represent afforestation. Three new plant functional types have been implemented in order to represent planted forests in temperate, tropical, and boreal regions. The parameters of these plantation types were estimated based on observations of stemwood carbon from real-world forest plantations. Using this new LPJmL version we present a global assessment of



- 1 potential carbon sequestration rates in forest plantations and compare these to rates achieved by
- 2 letting forests grow back naturally.

3 2 Methods

4 2.1 The LPJmL dynamic global vegetation model

5 LPJmL (Lund-Potsdam-Jena Managed Land) is a global process-based model simulating vegetation
6 dynamics and fluxes of carbon and water in the vegetation and soil of terrestrial ecosystems (Bondeau
7 et al., 2007; Schaphoff et al., 2013; Sitch et al., 2003), including agricultural land and biomass
8 plantations for bioenergy production (Beringer et al., 2011). The model runs primarily on a daily time
9 step, except for C allocation, vegetation dynamics, and disturbances for natural vegetation and
10 biomass plantations, which are resolved annually. Forcing consists of monthly climate variables (air
11 temperature, precipitation, cloud fraction, and number of wet days per month)—which are
12 interpolated to daily values (Gerten et al., 2004)—and annual atmospheric CO₂ concentrations. Using
13 a combination of plant physiological relations, generalized empirically established functions, and plant
14 trait parameters, LPJmL simulates processes such as photosynthesis, plant growth, maintenance and
15 regeneration losses, fire disturbance, soil moisture dynamics, runoff, evapotranspiration, irrigation
16 and vegetation structure (Schaphoff et al. 2013). Natural vegetation is represented as a number of
17 plant functional types (PFTs): aggregated vegetation classes representing variation in leaf-type
18 (broadleaf, needleleaf), phenology (summergreen, evergreen, raingreen), and climate preference
19 (boreal, temperate, tropical). Most model parameters related to vegetation are defined separately for
20 each PFT. The model simulates the occurrence of each PFT based on bioclimatic limits and competition
21 with other PFTs for resources. Agricultural ecosystems are handled in a separate module and are
22 represented by a range of crop functional types (Bondeau et al., 2007). Additionally, two woody and
23 one herbaceous PFTs are implemented to simulate short-rotation bioenergy plantations (Beringer et
24 al., 2011). Area fractions specifying allocation to different land-use types are part of the model input.
25 Finally, the model can simulate river discharge and surface water reservoirs, and several types of
26 irrigation. LPJmL has been coupled to the IMAGE integrated assessment model, serving as the land
27 surface component (Müller et al., 2016; Stehfest et al., 2014).

28 In all simulations for this study the model was forced by semi-constant monthly climate input,
29 representative for the period 1980–2010. This dataset was derived by repeating a cycle of detrended
30 time series for this period, taken from the CRU TS3.23 global gridded (0.5°×0.5° degrees) climate
31 dataset (Harris et al., 2014). For simplicity we chose to ignore the effect of atmospheric CO₂
32 concentration change at this stage, hence this variable was held fixed at the mean global value for



1 1980–2010 (362.4 ppmv). Fire disturbance was not considered. Further information on the model
2 input and configuration is given in subsequent sections.

3 2.2 Forest plantations

4 LPJmL was extended to represent forest plantations. Specifically, a new land-use type was added, as
5 well as three functional types to represent plantation trees in temperate, tropical, and boreal
6 plantations. These types—referred to as forest plantation functional types (FPFTs)—are derived from
7 the natural PFTs temperate broadleaved summergreen tree, tropical broadleaved evergreen tree, and
8 boreal needleleaved evergreen tree, respectively. The occurrence of the FPFTs is subject to the same
9 establishment and mortality rules used for natural PFTs. However, the bioclimatic limits are set such
10 that they do not overlap, hence co-occurrence of different FPFTs in a single grid cell is rare, occurring
11 only when climate fluctuates near a boundary between two types..

12 Structurally, the implementation largely follows that of the woody bioenergy plantations implemented
13 in LPJmL (Beringer et al., 2011), which in turn are based on equivalent natural PFTs. Contrary to
14 bioenergy trees, forest plantations are not automatically clear-cut after a fixed rotation period, but a
15 fraction of the plantation fraction may be harvested, specified as model input. However, for the
16 purpose of this study, harvest was set to zero. Forest plantation PFTs also differ from other PFTs with
17 regard to establishment of new trees. A fixed initial planting density (P_{init}) was introduced, which
18 determines the number of trees per unit area at planting. After planting, establishment of new trees
19 occurs similar to natural PFTs: at fixed maximum rate, downscaled according to an exponentially
20 declining function of foliar projective cover. Generally, stand density will decrease after plantation due
21 to self-thinning, implemented according to Reineke's rule (Reineke, 1933), which relates stem
22 diameter to crown area. When the area-fraction forest plantations in a given grid cell increases over
23 time, establishment is determined as a combination of P_{init} and the standard establishment rate,
24 weighted according to the old forest plantation fraction and the fraction added.




Figure 1. Location of the grid cells included in calibration simulations (100 per FPFT). The map shows simulated stemwood C (kgC m^{-2}) for a simulation with potential natural vegetation.

1 2.3 Calibration

2 2.3.1 General setup

3 To obtain realistic growth rates, we calibrated several FPFT-specific parameters, based on published
4 observed growth data for forest plantations. Ideally, calibration of dynamic vegetation models should
5 be performed using detailed observations for a given site. However, this requires a large amount of
6 data, both for model input and to compare to model output to assess performance. While much data
7 on growth of forest plantations has been published, the number of forest plantation sites for which
8 calibration data as well as data for model input are available for sufficiently long time periods is not
9 enough to derive globally applicable parameter sets. Therefore, we chose a different approach. Rather
10 than aiming to reproduce site-level observations, we calibrated the model in order to obtain desired
11 mean biome-level behavior, for each of the three FPFTs. For every iteration in the calibration, the
12 model was run for a selection of 100 grid cells from the spatial domain of the FPFT being calibrated.
13 Subsequently, model output for the relevant variables were aggregated over all grid cells and
14 compared to observed values to determine model performance.

15 Within the spatial domain for a given FPFT, many grid cells exist where growth is marginal due
16 unfavorable climate and/or soil properties. The observations used in the calibration are not
17 representative for these locations, since forest plantations from which data have been retrieved can
18 be assumed to represent locations where productivity is sufficient for economic profitability.
19 Therefore, rather than choosing grid cells randomly, the selection was limited to locations for which
20 LPJmL simulates relatively high productivity. This was done based on results from a 300-year simulation
21 with only natural vegetation, in the same setup as used in the calibration (see section 2.3). For each
22 FPFT, 100 cells were selected for which the simulated stemwood C storage of the corresponding
23 natural PFT (see section 2.2) exceeds the 70% percentile over the complete domain where this PFT is



1 dominant, i.e. has highest foliar projective cover (Fig. 1). During the calibration, LPJmL was run only for
2 these cells, with land-use type set to forest plantations.

3 2.3.2 Observations

4 *Target growth curves*

5 Time series of stand-level stemwood C were collected from various sources in the literature. For the
6 tropical FPFT we used data from Brown et al. (1986), who derived time series of stemwood biomass
7 for several species and species groups for tropical forest plantations. Data from natural poplar (*populus*
8 *x euramericana*) forests were taken from Cannell (1982) for the calibration of the temperate FPFT.
9 Finally, for the boreal FPFT we used data for Scots Pine (*Pinus Sylvestris*) plantations from Vanninen et
10 al. (1996). Outliers in the observations were removed using Hampel filtering (Pearson, 2002). The data
11 are depicted in Figure 2.

12 Since most forest plantations are grown for timber production, they are harvested approximately at
13 the optimal rotation length for maximum wood production, which is well before the trees reach
14 maturity. Hence, growth data for higher tree ages are scarce. Calibrating LPJmL against these
15 observations alone would result in excessive weight on the earlier part of the curve, leading to
16 unpredictable results for the later part. Therefore, we did not use the observations directly in the
17 LPJmL calibration but used them to derive growth curves representing the typical growing behavior of
18 productive plantations for each FPFT. We refer to these as the target growth curves (TGCs). The general
19 structure of the TGCs is given by the Chapman-Richards function, which is widely used to model forest
20 growth (e.g. Von Gadow and Hui, 1999). It defines the stemwood C (C_{SWC}) at time t as:

$$C_{SWC}(t) = C_{SWC,max}(1 - e^{-k t})^p, \quad (1)$$

21 where $C_{SWC,max}$ is the maximum C_{SWC} , k is the growth rate, and p is a shape parameter determining
22 the timing of maximum growth. These parameters were estimated using a Bayesian non-linear
23 regression approach. The scarcity of high-age observations was solved by constraining $C_{SWC,max}$ using
24 a prior distribution based on LPJmL output for the corresponding natural PFTs, from the 300-year
25 simulation used to select the calibration grid cells (see section 2.3.1). Specifically, for each FPFT we
26 used the mean simulated stemwood C of the last 10 simulation years, averaged over the 100
27 calibration cells as a representative value.

28 The parameters were estimated using MCMC sampling. The sample with highest posterior density,
29 together with the variances over the marginal posterior distributions, were used in the LPJmL
30 calibration. Further details are given in the supplemental text.



1 *Additional constraints*

2 Initial tests showed that parameter sets derived by calibration with the TGCs alone result in
 3 unrealistically high values of net primary production (NPP), leading to similarly high litter fluxes and
 4 soil carbon storage. This was traced to a higher carbon use efficiency (CUE)—the ratio of NPP to gross
 5 primary productivity—and a lower vegetation carbon turnover time (τ_{vegC} ; vegetation C to NPP ratio)
 6 compared to the natural PFT counterparts. Therefore, to assure realistic carbon fluxes and storage, we
 7 implemented additional constraints for these variables in the calibration. These constraints are based
 8 on LPJmL simulations for the natural PFT counterparts of the FPFTs, similar to the maximum stemwood
 9 C of the target growth curves (see Table 2 and Figure 4).

10 Additionally, it was found that certain parameter sets, while leading to acceptable mean results, cause
 11 simulated trees for certain cells to die-off repeatedly at regular intervals. In order to avoid this we
 12 modified the calibration such that a penalty was added to the cost function when this occurs.

Table 1. LPJmL parameters included in the calibration. Prior mode refers to the most probable value indicated by the prior distribution.

Parameter	Description	Units	Temperate		Tropical		Boreal	
			Prior mode	Estimate	Prior mode	Estimate	Prior mode	Estimate
α_a	Fraction of PAR assimilated at ecosystem level, relative to leaf level	-	0.5	0.61	0.5	0.60	0.5	0.53
g_{min}	Minimum canopy conductance	mm s ⁻¹	0.5	0.62	0.5	0.48	0.3	0.28
E_{max}	Maximum transpiration rate	mm d ⁻¹	5	5.5	7	11.7	5	6.07
r	Maintenance respiration coefficient	gC gN ⁻¹ d ⁻¹	1.2	1.5	0.2	0.16	1.2	1.2
k_{allom1}	Allometry parameter 1; relates crown area to stem diameter	-	100	126	100	166	110	86.4
k_{allom2}	Allometry parameter 2; relates tree height to stem diameter	-	40	53.9	40	41.1	40	36.5
k_{allom3}	Allometry parameter 3; relates tree height to stem diameter	-	0.67	1.02	0.67	0.84	0.67	1.2
lr_{max}	Leaf to root ratio under non-water stressed conditions	-	1	1.2	1	1.5	1	1.5
$C_{sapwood,sapl}$	Sapwood C of saplings	gC m ⁻²	1.2	1.02	1.2	1.2	1.2	1.4
LAI_{sapl}	Leaf area index of saplings	-	1.5	1.4	1.5	1.4	1.5	1.7
α_{leaf}	Leaf longevity	months	0.5	0.46	2	1.4	4	4.0
$\tau_{sapwood}$	Turnover time of sapwood	yr	20	20	20	46.3	20	15.7
$\tau_{leaf,root}$	Turnover time of leaves and roots	yr	1	1.3	2	1.8	4	4.7
P_{init}	Planting density	m ⁻²	0.15	0.15	0.15	0.15	0.15	0.17
k_{mort1}	Maximum mortality rate	yr ⁻¹	0.03	0.064	0.03	0.058	0.03	0.048



1 2.3.3 Parameter estimation

2 In the calibration 15 parameters were estimated, separately for each FPFT (Table 1). The calibration
 3 was performed on a transformed scale (logit for α_a ; log for all other parameters), in view of the lower
 4 bound at zero (and upper bound at 1 for α_a). We applied a Bayesian cost function, including
 5 informative prior distributions. Priors express belief about reasonable parameter values before the
 6 calibration in the form of probability distributions and help to avoid unrealistic values, particularly for
 7 parameters that have little influence on the relevant model output. The priors were chosen such that
 8 their central tendency reflects existing parameter values for the corresponding natural PFTs, with a
 9 relatively wide variance to avoid overly strong influence on the calibration. Full specification of the
 10 priors is given in supplemental text S1.

11 Similar to the parameters, all observations were transformed in the calibration (logit for CUE; log for
 12 all other observations). For the calibration simulations, LPJmL was started from zero vegetation and
 13 soil C and run for a period of 300 years, sufficient for the vegetation C to reach equilibrium with
 14 reasonable parameter values. LPJmL simulates heartwood and sapwood C pools, but does not
 15 distinguish between stem, branches, and coarse roots. For the purpose of the calibration, we assumed
 16 that all heartwood and 66% of the sapwood is located aboveground (Müller et al., 2016), and 84% of
 17 aboveground wood is located in the stem (which is representative for mature trees (Pretzsch, 2010)).

18 After simulation, the Chapman-Richards function was fitted to the time-series of simulated stemwood
 19 C for the 100 grid cells (using non-linear least squares) to derive FPFT-mean estimates of $C_{SWC,max}$, k ,
 20 and p based on LPJmL predictions. Carbon use efficiency and vegetation turnover time were
 21 determined for the last 10 years of the simulation, averaged over the 100 grid cells. The observations
 22 were subsequently compared to the corresponding observations to determine log-likelihood, and
 23 combined with log prior density to determine the overall cost $\mathcal{C}(\theta)$ for the given parameter set θ .
 24 Further details are given in supplemental text S2.

Table 2. Observations and corresponding fits for the 100 included grid cells included in the calibration.
 Observed values correspond to the mode of the likelihood function.

Symbol	Description	Units	Temperate		Tropical		Boreal	
			Obs.	Fit	Obs.	Fit	Obs.	Fit
$C_{SWC,max}$	Growth curve parameter; maximum stemwood C	kgC m ⁻²	6.77	6.45	15.62	15.74	7.45	7.56
k	Parameter of growth curve	yr ⁻¹	0.197	0.0420	0.0566	0.0301	0.0500	0.0257
p	Parameter of growth curve	-	3.91	3.37	1.59	1.69	4.28	4.64
CUE	Carbon use efficiency; NPP to GPP ratio	-	0.380	0.342	0.458	0.448	0.460	0.427
τ_{vegC}	Vegetation C turnover time; Vegetation C to NPP ratio	yr	16.86	18.37	21.92	17.99	22.27	19.52



1 The optimal parameter set (with minimal value of C) was derived using the genoud algorithm (Mebane
2 Jr. and Sekhon, 2011) which combines a genetic algorithm with a gradient search approach. This
3 algorithm has previously been applied to calibrate LPJmL (Forkel et al., 2014). Additional description is
4 given in supplemental text S2.

5 2.4 Global simulations

6 After calibration, several global simulations were performed. First, in order to assess sequestration
7 potential of afforestation, a simulation was run in the same setup as used for the calibration, i.e.
8 starting with zero vegetation and soil C and with land fully allocated to forest plantations and running
9 for 300 years so that vegetation C pool can reach equilibrium. Additionally, a simulation with land fully
10 allocated to natural vegetation was performed, to compare natural regrowth and afforestation as land-
11 based mitigation options.

12 Second, we applied the model for an ambitious scenario of large-scale afforestation, assuming that
13 from 2015 onwards approximately 14% of global managed land is (corresponding to 650 Mha or 4.5%
14 of global land surface) gradually replaced by forest plantations over the course of 85 years. This
15 afforestation area is in line with the average land area used for land-based mitigation (both bio-energy
16 and afforestation) in 1.5 degree mitigation scenarios in Integrated Assessment Models (Doelman et
17 al., in review; Rogelj et al., 2018). To bring soil C to reasonable values, the simulation was initialized by
18 two spin-up phases: 1) a 1000-year phase with natural vegetation only until 1901, and 2) a phase from
19 1900 to 2015 with transient cropland and pasture fractions, based on the HYDE dataset (Klein
20 Goldewijk et al., 2010). From 2015 forest plantation area was increased and crop and pasture area
21 was, balancing each other so that total area of managed (i.e. non-natural) land remained constant.
22 From 2100 the simulation was continued for another 50 years with constant land-use. For this analysis,
23 two complementary simulations were performed. First, a simulation where fractions of natural
24 vegetation were increased, instead of forest plantations, and second, a “baseline” simulation where
25 land-use fractions were held constant in time from 2015. Supplemental Figures S2 and S3 depict the
26 development of land-use fractions for the three scenarios.

27 3 Results

28 3.1 Target growth curves

29 Figure 2 depicts the stemwood C observations for the three FPFTs, LPJmL simulations for the
30 corresponding natural PFTs, and the target growth curves (TGCs) resulting from the fitting procedure.
31 The values of the maximum stemwood C ($C_{SWC,max}$), growth rate (k), and shape parameter (p) and
32 their marginal variances are given in Table 2 (see also Figure S1). As expected, the observations show
33 substantially higher growth rates than simulated for the natural PFTs, as well as an earlier timing of



- 1 maximum growth. The TGCs represent a compromise between the observations and the $C_{\text{SWC,max}}$ for
- 2 natural PFTs, predicted by LPJmL: the initial high growth rate is representative for the observations,
- 3 while $C_{\text{SWC,max}}$ is closer to that of the simulated natural PFTs and notably lower than the level
- 4 indicated by the observations.
- 5 The tropical FPFT has substantially higher $C_{\text{SWC,max}}$, approximately twice as high as the other two
- 6 FPFTs. With respect to the relative growth rate k , however, the tropical FPFT is comparable to the
- 7 boreal FPFT. The temperate FPFT approaches its maximum stemwood approximately four times faster.
- 8 The boreal TGC has the highest value of p , resulting in a later timing of maximum growth.




Figure 2. Target growth curves for stemwood C, associated observations and LPJmL output for natural vegetation in the cells selected for calibration.




Figure 3. Prior distributions and estimated final values of the FPFT parameters estimated in the calibration. The boxplots indicate the 5% and 95% percentiles (whiskers) the median (red line) and 25% and 75% percentiles (box) or the priors. The final parameter estimate is indicated by the asterisk (*). See Table 1 for explanation of the parameters.




Figure 4. Ranges of the observations used in the calibration and LPJmL estimates after calibration. The boxplots indicate the 5% and 95% percentiles (whiskers) the median (red line) and 25% and 75% percentiles (box) of the likelihood function. The final fitted value is indicated by the asterisk (*). $C_{SWC,max}$: maximum stemwood C, k : growth rate; p : shape factor, CUE: carbon use efficiency (NPP to GPP ratio); τ_{vegC} : vegetation C turnover time (vegetation C to NPP ratio).



1 3.2 LPJmL calibration

2 The parameter estimates resulting from the calibration are shown in Figure 3, together with the range
3 of the prior distributions. Most estimates are within interquartile range of the priors, but for several
4 parameters the calibration resulted in relatively strong changes, in particular $k_{\text{mort}1}$, which controls
5 mortality due to low growth efficiency. Specifically for the tropical FPFT, the estimates also clearly
6 deviate from the prior for E_{max} , $k_{\text{allom}1}$, $l_{r_{\text{max}}}$, and τ_{sapwood} .

7 The ranges of the observed variables are depicted in Figure 4, together with the LPJmL predictions for
8 the calibration grid cells, based on the optimized parameter sets. The parameter $C_{\text{SWC,max}}$ is fit well
9 by the model for all FPFTs. However, k is clearly underestimated for all three FPFTs, compared to the
10 observed ranges. This affects the simulated growth curve for stemwood C, as shown in Figure 5. In the
11 LPJmL simulation, vegetation needs a longer time to reach its maximum stemwood biomass than the
12 target growth curve. Nevertheless, the growth rate based on the optimized parameters represents a
13 substantial improvement compared to the natural PFT counterparts. The carbon use efficiency (CUE)
14 and vegetation turnover time (τ_{vegC}) are also reasonably well fitted.




Figure 5. Predicted stemwood C for 100 calibration grid cells of each FPFT based on the optimal parameter sets. Note the different scales of the y axes.




Figure 6. Spatial distribution of the different forest plantation functional types resulting from the bioclimatic limits. In marginal regions no trees are simulated, but grass may be present.




Figure 7. Global total ecosystem C over time for simulations with global forest plantations or global natural vegetation. (a) vegetation C storage per biome for the global forest plantation simulation only; (b) C storage per compartment for both simulations.

1 3.3 Global simulations

2 3.3.1 Global afforestation and natural regrowth

3 Figure 6 depicts the predicted spatial distribution of forest plantation functional types for a global
 4 simulation experiment with land fully allocated to forest plantations. The total area for the temperate,
 5 tropical and boreal plantation types is 2,472 Mha (10^{10} m²), 6,242 Mha, and 3,094 Mha, respectively,
 6 corresponding to 17%, 43%, and 22%, of global land surface. In 2,579 Mha (18%) no tree growth is
 7 simulated due to a too cold or too dry climate. Note that in many non-marginal regions tree growth
 8 may still be very low due to unfavorable conditions—the depicted distribution simply results from the
 9 bioclimatic limits set in the model. The distribution the FPFTs corresponds roughly to zones C, A, and
 10 D of the first level of the Köppen climate classification (Peel et al., 2007). Since there is no type for
 11 forest plantations in arid climates, the three FPFTs extend also into desert regions.

12 The global vegetation C stock over time is depicted in Figure 7a (cf also Figure S2). Tropical plantations
 13 contribute most to C storage due to their larger area and higher productivity. Comparison with the
 14 simulation where all land is allocated to natural vegetation shows considerably faster C uptake for
 15 forest plantations (Figure 7b), with a maximum difference of 308 PgC (193%), after 54 years. After 300




Figure 8. Mean ecosystem sequestration rate (Mean annual increment, MAI), determined as total C storage divided by time since start for LPJmL simulations with only forest plantations or only natural vegetation. (a) global total versus time; (b) difference between (afforestation minus natural regrowth) after 50 years.

1 years, global vegetation C is 102 PgC higher (112%) for afforestation simulation. Soil and litter C storage
2 is also proportionally higher for forest plantations. Note that the soil and litter C uptake rate is
3 extremely high due to the fact that the simulation was started with zero C. In reality soil C will already
4 be present before land-use change and uptake will be much slower, possibly even negative, depending
5 on previous land-use.

6 The potential for C uptake is illustrated by the mean annual increment (MAI) of vegetation C since the
7 start of plantation (Figure 8). There are remarkable differences between the two simulations. After an
8 initial similar increase, MAI sharply drops after approximately 10 years for natural regrowth, while for
9 afforestation MAI keeps rising until approximately 30 years. The behavior for natural regrowth can be
10 explained by vegetation succession, leading to a shift from grasses to trees. This succession does not
11 occur for forest plantations, where trees start growing immediately, resulting in a substantially higher
12 MAI in the early part of the simulation. From spatial differences in MAI after 50 years it is evident that
13 tropical regions contribute most to this difference.

14 3.3.2 Transient afforestation and natural regrowth

15 Figure 9 depicts results of the global simulation scenarios with gradual increase in forest, applying
16 either afforestation or natural regrowth. Since changes in C storage—particularly for the soil—result
17 also from land-use changes before




Figure 9 Results of the simulations for gradual afforestation and natural regrowth. Both graphs show differences relative to the baseline simulation with constant land-use. (a) global C storage; (b) global C sequestration rate, smoothed using a 10-year moving average window.

1 2015, we focus on the difference in global C stocks compared to the baseline simulation with constant
2 land-use from 2015. Until 2015 all three simulations have very similar results, but small differences
3 arise from the stochastic generation of daily precipitation. Gradual afforestation of 650 Mha land
4 between 2015 and 2100 results in 19, 48, and 75 PgC additional C storage by 2065, 2100, and 2150,
5 respectively, versus 16, 37, and 61 PgC for natural regrowth. Most of the difference between the two
6 simulations is due to vegetation C, but from 2100 the difference for soil C grows and would ultimately
7 dominate, had the simulation been continued after 2150. Global C sequestration rate peaks between
8 2090 and 2100 at approximately 0.91 and 0.68 PgC yr⁻¹ for afforestation and natural regrowth, with
9 average rates of 0.25 and 0.19 PgC yr⁻¹ until 2100. Around 2130 global sequestration rates are higher
10 for the natural regrowth simulation because land-use remains constant from 2100, allowing natural
11 ecosystems to “catch up”.

12 4 Discussion

13 4.1 LPJmL calibration

14 4.1.1 Parameters changes

15 Compared to the prior distributions—which are largely based on values for corresponding natural
16 PFTs—the calibration resulted in a substantial shift for several parameters. We will discuss the more
17 notable changes. First, k_{mort1} is substantially higher compared to the prior mode for all FPFTs. This
18 parameter controls tree mortality related to low growth efficiency, which is defined as the ratio of the
19 annual net biomass increment to leaf area. A high value of k_{mort1} results in higher mortality under
20 unfavorable conditions. The increase of this parameter can be explained by the fact that the target
21 growth curves have substantially higher growth rates than the natural PFT equivalents, while maximum
22 biomass is approximately the same. The apparent conflict between these two constraints can in part



1 be resolved by increasing first-order mortality. A higher value for k_{mort1} for forest plantation trees is
2 not necessarily unrealistic since it is likely that fast-growing tree species have low tolerance for low
3 growth conditions (Pacala et al., 1996).

4 The parameter k_{allom3} , which relates tree stem diameter to tree height, has also substantially
5 increased for all FPFTs. Higher values of this parameter mean higher trees for the same diameter,
6 resulting in higher maximum biomass per tree. Again, this is in agreement with field observations,
7 which have shown a positive relationship between tree growth rate and this parameter (Martinez Cano
8 et al., 2018)

9 The maximum leaf-to-root mass ratio, l_{rmax} , is also high compared to the prior, particularly for the
10 tropical and boreal FPFTs. This causes higher allocation of C to leaves, compared to roots, which
11 positively affects growth rate via leaf area index and absorbed photosynthetically active radiation.
12 Conversely, in LPJmL, lower root biomass does not reduce growth since there is no link between root
13 biomass and water uptake. Hence, higher values of l_{rmax} unequivocally lead to higher productivity in
14 the model.

15 For the tropical FPFT, E_{max} , τ_{sapwood} , and k_{allom1} have increased. E_{max} is the maximum water
16 transport capacity and controls the transpiration rate. τ_{sapwood} is the turnover time of sapwood—
17 higher values result in more sapwood biomass, which allows for larger leaf area. Finally, k_{allom1} relates
18 crown area to stem diameter. The generally stronger shifts for the tropical FPFT compared to
19 temperate and boreal is explained by the lower uncertainties of parameters k and p of the target
20 growth curve (Figure 4, S1), which is in turn caused by the larger number of stemwood C observations
21 (Figure 2).

22 4.1.2 Fit to observations

23 The calibration resulted in good fits with respect to most observations, with the exception of the
24 growth rate parameter of the target growth curves. Despite substantial improvement compared to the
25 corresponding natural PFTs, this parameter is underestimated for all three FPFTs. As a result, predicted
26 initial C uptake rates are lower than implied by the stemwood C observations, possibly underestimating
27 the potential efficacy of forest plantations for climate mitigation.

28 As discussed in section 2.3.2, we incorporated data into the calibration to constrain NPP to GPP ratio
29 and vegetation C turnover time to values similar to that of the corresponding natural PFTs. Earlier
30 calibrations, in which these constraints were not included, yielded a substantially better fit to the
31 growth rate but with unrealistic litter fluxes, which points to a trade-off between the fit to these
32 observations. From a mass-balance perspective this result is explicable: fast growth requires high NPP,
33 which will result in high litter fluxes once vegetation reaches equilibrium biomass. This is exacerbated



1 by the fact that we constrained maximum stemwood C ($C_{SWC,max}$) at levels close to that of the
2 corresponding natural PFTs. The fact that fast growth results in very high litter fluxes when trees reach
3 equilibrium relates to the fact that LPJmL does not represent certain mechanisms that lead to declining
4 productivity with age (Zaehle et al., 2006). In reality, NPP reduction with age has been frequently
5 observed (Ryan et al., 1997). Multiple reasons for this phenomenon have been proposed, but the
6 leading hypothesis is that hydraulic resistance increases with tree height due to longer distance
7 between soil and leaves (Ryan and Yoder, 1997). This result in a lower photosynthesis rates and gross
8 productivity (GPP). Since LPJmL does not include such mechanisms, it is mostly representative for
9 mature forests. Incorporation of a more realistic representation of age-dependence of forest growth
10 rate is likely to improve fit to observations (Zaehle et al., 2006).

11 4.2 Global simulations

12 4.2.1 Global afforestation versus natural regrowth

13 Despite the underestimates growth rates, our results show C can be sequestered substantially faster
14 by forest plantations compared to natural regrowth (Figure 7), particularly in the first 50 years
15 following land conversion. The largest potential for plantations lies in tropical regions, which is not
16 surprising, given that the maximum biomass of tropical FPFTs is more than twice as high compared to
17 the temperate and boreal FPFTs. In addition to faster C sequestration, LPJmL also predicts a 12% higher
18 equilibrium global vegetation C pool for forest plantations, despite the fact that the FPFTs were
19 calibrated to produce a value of $C_{SWC,max}$ comparable to that of natural equivalent PFTs (Figure 5).
20 This contradiction is in part explained by a larger productivity of forest plantations in less productive
21 regions, which were not included in the calibration grid cells. Another reason is a larger spatial extent
22 of the tropical FPFT compared to that of the natural tropical PFTs.

23 Combined soil and litter C is also higher for forest plantations after 300 years (Figure 7b), but its
24 proportion to total ecosystem C (60%) is globally almost identical to that of the natural vegetation,
25 owing to the constraints on NPP to GPP ratio and vegetation C turnover time included in the calibration
26 (see section 2.3.2). It is difficult to compare these results to observations for real-world plantations
27 since studies on this topic have generally compared natural forests to tree plantations for production
28 of wood or other products, where the effects of harvest and other management on soil C are likely
29 considerable (Guo and Gifford, 2002; van Straaten et al., 2015). Such effects are not relevant for
30 plantations intended for C sequestration.

31 4.2.2 Gradual afforestation versus natural regrowth

32 According to our projections, gradual conversion of 650 Mha managed land to natural forest between
33 2015 and 2100 results in additional C uptake of 16 and 37 PgC by 2065 and 2100, respectively. If these
34 lands are converted to forest plantations, the estimated C uptake is 19 and 48 PgC, i.e. 19% and 30%



1 higher. These should be seen as a conservative estimates, in view of the underestimated growth rates
2 resulting from the calibration. To put these numbers into perspective, we compare them to results of
3 Gasser et al. (2015), who estimated the negative emissions needed to limit global warming to 2°C for
4 a range of scenarios in which both the start time and the rate of reduction of greenhouse gasses varied.
5 In their most favorable scenario (energy and industry emission reduction starting in 2015 at a rate of
6 5% per year) they estimated an average cumulative negative emission of 25–100 PgC is needed by
7 2100, compared to 450–800 PgC in the most unfavorable scenario (energy and industry emission
8 reduction starting in 2030 at 1% per year). Hence, large-scale forest plantations can offer a substantial
9 contribution to climate mitigation but will likely not be sufficient.

10 4.3 Comparison to previous work

11 The results of the simulations for transient afforestation and natural regrowth compare well to results
12 of previous studies on potential C sequestration rates of forest plantations and natural regrowth. For
13 example, using the IMAGE integrated assessment model, van Minnen et al. (2008) performed a
14 simulation experiment based on the IPCC SRES A1B scenario where 831 Mha agricultural land is
15 converted to permanent forest plantations between 2000 and 2100, taking into consideration land
16 demand for food production and other uses. They estimated an additional 93 PgC can be sequestered,
17 but mostly after 2050, when land becomes gradually available due to decreasing population and
18 increasing agricultural efficiency.

19 Humpenöder et al. (2014) presented a much more ambitious afforestation scenario, in which 2773
20 Mha land is converted to forest plantations. The authors used maximum C storage for natural
21 vegetation predicted by LPJmL, but corrected sequestration rates using stylized growth curves for
22 plantations in different climate regions. They estimate an additional C uptake of 192 PgC after 80 years.
23 Roughly converting our estimate to the same land area yields a similar result (205 PgC). This similarity
24 is not surprising, given that we used the same model, and our FPFTs were calibrated to produce the
25 same maximum biomass as the natural PFT equivalents.

26 Potential sequestration rates by natural regrowth were studied by Krause et al. (2017), using the
27 dynamic global vegetation model LPJ-GUESS. In two scenarios, derived by IMAGE and the agricultural
28 land-use model MAgPIE, 1119 and 914 Mha were converted to natural lands, resulting in a predicted
29 additional C uptake of 76 and 55 PgC, respectively, between 2000 and 2099. This compares well with
30 our estimates for natural regrowth.

31 4.4 Model limitations

32 In our implementation of planted forests the diversity of plantation tree species is reduced to three
33 functional types with fixed properties. While the functional diversity of plantation tree species is not



1 as vast as that of natural forests—especially in the context of C sequestration—the predictions would
2 likely improve from implementation of additional FPFTs, particularly for the tropical biome. The model
3 currently predicts a relatively large C storage for dry tropical zone compared to natural regrowth,
4 which may not be fully realistic, given water limitations. Addition of a dry tropical FPFT would allow for
5 a more accurate assessment of C sequestration in these regions.

6 This study does not consider the effects of climate change and CO₂ concentration on productivity of
7 forest plantations. Although there is still considerable uncertainty regarding this topic, accounting for
8 CO₂ fertilization will likely increase the C sequestration potential (Schimel et al., 2015) in both natural
9 and managed forests. However, in order to properly assess this, it is important to take into account
10 nutrient limitation to productivity as well (Norby et al., 2010).

11 We also did not consider possible management options that may improve C uptake rates. In particular,
12 regular thinning can result in substantially higher C uptake rates (van Minnen et al., 2008). The model
13 supports harvesting, but this feature was not used in this study. However, continual thinning would
14 result in export of nutrients from the ecosystem which would ultimately slow down growth rates,
15 unless plantations are fertilized. Thus, representing regular harvest in LPJmL would also require
16 representation of nutrient limitation.

17 4.5 Considerations beyond C uptake

18 Evaluation of afforestation and natural regrowth as strategies for climate change mitigation involves a
19 range of considerations other than carbon sequestration. First, converting agricultural land to forest
20 involves a number of costs. For both natural and planted forests this includes price for acquiring land,
21 while specifically for the latter costs related to establishing and maintaining the plantation are relevant
22 (e.g. land preparation, planting of seedlings). The costs per unit C sequestered will rise with increasing
23 area of (planted) forest, mainly due to competition for land (Doelman et al., in review).

24 Second, the positive effects of carbon uptake of changing land-cover to forest, can be offset due
25 biophysical changes in the surface energy budget, related to changes in albedo, evapotranspiration,
26 and surface roughness (Perugini et al., 2017). This may result in a net warming effect, regionally, and
27 possibly globally, depending on the extent of land-cover change.

28 Third, the reduction of cropland and pasture might also have a negative impact on food security due
29 to increased competition for land (Hasegawa et al., 2018). In order to maintain food production for the
30 growing population, strong intensification of the agricultural sector would be required. Locally, this
31 will result in a range of negative effects on the environment, due to higher application of fertilizers and
32 plant protection products, as well as water extraction for irrigation (Smith et al., 2013). Furthermore,



1 in terms of climate change mitigation, agricultural intensification will likely partially offset the benefits
2 of afforestation and regrowth, e.g. due to higher N₂O emissions from fertilizers (Burney et al., 2010).
3 Finally, biodiversity is a particularly important aspect to consider, given that plantation forests have
4 usually substantially lower species richness than primary or secondary forests (Barlow et al., 2007). A
5 more balanced solution may be a compromise between biodiversity and C sequestration by
6 establishing a mixture of native and plantation species, or plantation forest with a native undergrowth
7 (Barlow et al., 2007; Bremer and Farley, 2010).

8 5 Conclusions

9 To our knowledge, the extension of LPJmL presented here represents the first model of forest
10 plantations for C sequestration as part of a DGVM for global-scale applications. Although calibration
11 of the model still resulted in underestimated growth rates compared to observations of stemwood C,
12 this represents an improvement over previous approaches. According to our simulations, conversion
13 of 650 Mha of land to forest over 85 years results in an additional C uptake of 48 PgC for forest
14 plantation, versus 37 PgC for natural regrowth, with greatest potential in the tropics. We conclude that
15 large scale afforestation can offer a substantial contribution to C uptake, particularly in a time scale of
16 approximately 50–100 years. Evaluating afforestation as a strategy for climate change mitigation
17 requires consideration of all relevant aspects in a comprehensive assessment. Our model can
18 contribute to such an evaluation by providing improved estimates of C uptake rates.

19 Acknowledgements

20 We gratefully acknowledge Matthias Forkel for advice on using the Genoud algorithm to calibrate
21 LPJmL.

22

23



1 References

- 2 Barlow, J., Gardner, T. A., Araujo, I. S., Avila-Pires, T. C., Bonaldo, A. B., Costa, J. E., Esposito, M. C.,
3 Ferreira, L. V., Hawes, J., Hernandez, M. I. M., Hoogmoed, M. S., Leite, R. N., Lo-Man-Hung, N. F.,
4 Malcolm, J. R., Martins, M. B., Mestre, L. A. M., Miranda-Santos, R., Nunes-Gutjahr, A. L., Overal, W. L.,
5 Parry, L., Peters, S. L., Ribeiro-Junior, M. A., da Silva, M. N. F., da Silva Motta, C. and Peres, C. A.:
6 Quantifying the biodiversity value of tropical primary, secondary, and plantation forests, *Proc. Natl.*
7 *Acad. Sci.*, 104(47), 18555–18560, doi:10.1073/pnas.0703333104, 2007.
- 8 Beringer, T. I. M., Lucht, W. and Schaphoff, S.: Bioenergy production potential of global biomass
9 plantations under environmental and agricultural constraints, *GCB Bioenergy*, 3(4), 299–312,
10 doi:10.1111/j.1757-1707.2010.01088.x, 2011.
- 11 Bondeau, A., Smith, P. C., Zaehle, S., Schaphoff, S., Lucht, W., Cramer, W., Gerten, D., Lotze-Campen,
12 H., Mueller, C., Reichstein, M. and Smith, B.: Modelling the role of agriculture for the 20th century
13 global terrestrial carbon balance, *Glob. Change Biol.*, 13(3), 679–706, doi:10.1111/j.1365-
14 2486.2006.01305.x, 2007.
- 15 Bremer, L. L. and Farley, K. A.: Does plantation forestry restore biodiversity or create green deserts? A
16 synthesis of the effects of land-use transitions on plant species richness, *Biodivers. Conserv.*, 19(14),
17 3893–3915, doi:10.1007/s10531-010-9936-4, 2010.
- 18 Brown, S., Lugo, A. E. and Chapman, J.: Biomass of tropical tree plantations and its implications for the
19 global carbon budget, *Can. J. For. Res.-Rev. Can. Rech. For.*, 16(2), 390–394, doi:10.1139/x86-067,
20 1986.
- 21 Burney, J. A., Davis, S. J. and Lobell, D. B.: Greenhouse gas mitigation by agricultural intensification,
22 *Proc. Natl. Acad. Sci.*, 107(26), 12052–12057, doi:10.1073/pnas.0914216107, 2010.
- 23 Cannell, M. G. R.: *World forest biomass and primary production data*, Academic Press., 1982.
- 24 Doelman, J. C., Stehfest, E., van Vuuren, D. P., Tabeau, A., Hof, A. F., Braakhekke, M. C., Gernaat, D. E.
25 H. J., van den Berg, M., van Zeist, W., Daioglou, V., van Meijl, H. and Lucas, P.: Estimating afforestation
26 potentials and possible risks to food security, *Glob. Change Biol.*, in review.
- 27 Doelman, J. C., Stehfest, E., Tabeau, A. and van Meijl, H.: Making the Paris agreement climate targets
28 consistent with Food Security objectives, *Glob. Food Secur.*, in review.
- 29 Forkel, M., Carvalhais, N., Schaphoff, S., v. Bloh, W., Migliavacca, M., Thurner, M. and Thonicke, K.:
30 Identifying environmental controls on vegetation greenness phenology through model–data
31 integration, *Biogeosciences*, 11(23), 7025–7050, doi:10.5194/bg-11-7025-2014, 2014.
- 32 Gasser, T., Guivarch, C., Tachiiri, K., Jones, C. D. and Ciais, P.: Negative emissions physically needed to
33 keep global warming below 2 °C, *Nat. Commun.*, 6(1), doi:10.1038/ncomms8958, 2015.
- 34 Gerten, D., Schaphoff, S., Haberlandt, U., Lucht, W. and Sitch, S.: Terrestrial vegetation and water
35 balance—hydrological evaluation of a dynamic global vegetation model, *J. Hydrol.*, 286(1–4), 249–270,
36 doi:10.1016/j.jhydrol.2003.09.029, 2004.
- 37 Gladstone, W. T. and Thomas Ledig, F.: Reducing pressure on natural forests through high-yield
38 forestry, *For. Ecol. Manag.*, 35(1–2), 69–78, doi:10.1016/0378-1127(90)90232-Z, 1990.



- 1 Guo, L. B. and Gifford, R. M.: Soil carbon stocks and land use change: a meta analysis, *Glob. Change*
2 *Biol.*, 8(4), 345–360, doi:10.1046/j.1354-1013.2002.00486.x, 2002.
- 3 Harris, I., Jones, P. D., Osborn, T. J. and Lister, D. H.: Updated high-resolution grids of monthly climatic
4 observations - the CRU TS3.10 Dataset: UPDATED HIGH-RESOLUTION GRIDS OF MONTHLY CLIMATIC
5 OBSERVATIONS, *Int. J. Climatol.*, 34(3), 623–642, doi:10.1002/joc.3711, 2014.
- 6 Hasegawa, T., Fujimori, S., Havlík, P., Valin, H., Bodirsky, B. L., Doelman, J. C., Fellmann, T., Kyle, P.,
7 Koopman, J. F. L., Lotze-Campen, H., Mason-D’Croz, D., Ochi, Y., Pérez Domínguez, I., Stehfest, E.,
8 Sulser, T. B., Tabeau, A., Takahashi, K., Takakura, J., van Meijl, H., van Zeist, W.-J., Wiebe, K. and Witzke,
9 P.: Risk of increased food insecurity under stringent global climate change mitigation policy, *Nat. Clim.*
10 *Change*, 8(8), 699–703, doi:10.1038/s41558-018-0230-x, 2018.
- 11 Humpenoder, F., Popp, A., Dietrich, J. P., Klein, D., Lotze-Campen, H., Bonsch, M., Bodirsky, B. L.,
12 Weindl, I., Stevanovic, M. and Muller, C.: Investigating afforestation and bioenergy CCS as climate
13 change mitigation strategies, *Environ. Res. Lett.*, 9(6), doi:10.1088/1748-9326/9/6/064029, 2014.
- 14 Klein Goldewijk, K., Beusen, A. and Janssen, P.: Long-term dynamic modeling of global population and
15 built-up area in a spatially explicit way: HYDE 3.1, *The Holocene*, 20(4), 565–573,
16 doi:10.1177/0959683609356587, 2010.
- 17 Krause, A., Pugh, T. A. M., Bayer, A. D., Doelman, J. C., Humpenöder, F., Anthoni, P., Olin, S., Bodirsky,
18 B. L., Popp, A., Stehfest, E. and Arneith, A.: Global consequences of afforestation and bioenergy
19 cultivation on ecosystem service indicators, *Biogeosciences*, 14(21), 4829–4850, doi:10.5194/bg-14-
20 4829-2017, 2017.
- 21 Martinez Cano, I., Muller-Landau, H. C., Wright, S. J., Bohlman, S. A. and Pacala, S. W.: Interspecific
22 variation in tropical tree height and crown allometries in relation to life history traits, *Biogeosciences*
23 *Discuss.*, 1–25, doi:10.5194/bg-2018-314, 2018.
- 24 Mebane Jr., W. R. and Sekhon, J. S.: Genetic Optimization Using Derivatives: The rgenoud Package for
25 R, *J. Stat. Softw.* Vol 1 Issue 11 2011, doi:10.18637/jss.v042.i11, 2011.
- 26 van Minnen, J. G., Strengers, B. J., Eickhout, B., Swart, R. J. and Leemans, R.: Quantifying the
27 effectiveness of climate change mitigation through forest plantations and carbon sequestration with
28 an integrated land-use model, *Carbon Balance Manag.*, 3(1), 3, 2008.
- 29 Müller, C., Stehfest, E., van Minnen, J. G., Strengers, B., von Bloh, W., Beusen, A. H. W., Schaphoff, S.,
30 Kram, T. and Lucht, W.: Drivers and patterns of land biosphere carbon balance reversal, *Environ. Res.*
31 *Lett.*, 11(4), 044002, doi:10.1088/1748-9326/11/4/044002, 2016.
- 32 Norby, R. J., Warren, J. M., Iversen, C. M., Medlyn, B. E. and McMurtrie, R. E.: CO₂ enhancement of
33 forest productivity constrained by limited nitrogen availability, *Proc. Natl. Acad. Sci.*, 107(45), 19368–
34 19373, doi:10.1073/pnas.1006463107, 2010.
- 35 Pacala, S. W., Canham, C. D., Saponara, J., Silander, J. A., Kobe, R. K. and Ribbens, E.: Forest Models
36 Defined by Field Measurements: Estimation, Error Analysis and Dynamics, *Ecol. Monogr.*, 66(1), 1–43,
37 doi:10.2307/2963479, 1996.
- 38 Paquette, A. and Messier, C.: The role of plantations in managing the world’s forests in the
39 Anthropocene, *Front. Ecol. Environ.*, 8(1), 27–34, doi:10.1890/080116, 2010.
- 40 Pearson, R. K.: Outliers in process modeling and identification, *IEEE Trans. Control Syst. Technol.*, 10(1),
41 55–63, doi:10.1109/87.974338, 2002.



- 1 Peel, M. C., Finlayson, B. L. and McMahon, T. A.: Updated world map of the Köppen-Geiger climate
2 classification, *Hydrol. Earth Syst. Sci.*, 11(5), 1633–1644, doi:10.5194/hess-11-1633-2007, 2007.
- 3 Perugini, L., Caporaso, L., Marconi, S., Cescatti, A., Quesada, B., de Noblet-Ducoudré, N., House, J. I.
4 and Arneth, A.: Biophysical effects on temperature and precipitation due to land cover change,
5 *Environ. Res. Lett.*, 12(5), 053002, doi:10.1088/1748-9326/aa6b3f, 2017.
- 6 Pretzsch, H.: *Forest Dynamics, Growth and Yield*, Springer Berlin Heidelberg, Berlin, Heidelberg., 2010.
- 7 Reineke, L. H.: Perfecting a stand-density index for even-aged forests, *J. Agric. Res.*, 46, 0627–0638,
8 1933.
- 9 Rogelj, J., Popp, A., Calvin, K. V., Luderer, G., Emmerling, J., Gernaat, D., Fujimori, S., Strefler, J.,
10 Hasegawa, T., Marangoni, G., Krey, V., Kriegler, E., Riahi, K., van Vuuren, D. P., Doelman, J., Drouet, L.,
11 Edmonds, J., Fricko, O., Harmsen, M., Havlík, P., Humpenöder, F., Stehfest, E. and Tavoni, M.: Scenarios
12 towards limiting global mean temperature increase below 1.5 °C, *Nat. Clim. Change*, 8(4), 325–332,
13 doi:10.1038/s41558-018-0091-3, 2018.
- 14 Ryan, M. G. and Yoder, B. J.: Hydraulic Limits to Tree Height and Tree Growth, *BioScience*, 47(4), 235–
15 242, doi:10.2307/1313077, 1997.
- 16 Ryan, M. G., Binkley, D. and Fownes, J. H.: Age-Related Decline in Forest Productivity: Pattern and
17 Process, in *Advances in Ecological Research*, vol. 27, pp. 213–262, Elsevier., 1997.
- 18 Schaphoff, S., Heyder, U., Ostberg, S., Gerten, D., Heinke, J. and Lucht, W.: Contribution of permafrost
19 soils to the global carbon budget, *Environ. Res. Lett.*, 8(1), 014026, doi:10.1088/1748-
20 9326/8/1/014026, 2013.
- 21 Schimel, D., Stephens, B. B. and Fisher, J. B.: Effect of increasing CO₂ on the terrestrial carbon cycle,
22 *Proc. Natl. Acad. Sci.*, 112(2), 436–441, doi:10.1073/pnas.1407302112, 2015.
- 23 Sitch, S., Smith, B., Prentice, I. C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J. O., Levis, S., Lucht,
24 W., Sykes, M. T., Thonicke, K. and Venevsky, S.: Evaluation of ecosystem dynamics, plant geography
25 and terrestrial carbon cycling in the LPJ dynamic global vegetation model, *Glob. Change Biol.*, 9(2),
26 161–185, doi:10.1046/j.1365-2486.2003.00569.x, 2003.
- 27 Smith, P., Haberl, H., Popp, A., Erb, K., Lauk, C., Harper, R., Tubiello, F. N., de Siqueira Pinto, A., Jafari,
28 M., Sohi, S., Masera, O., Böttcher, H., Berndes, G., Bustamante, M., Ahammad, H., Clark, H., Dong, H.,
29 Elsiddig, E. A., Mbow, C., Ravindranath, N. H., Rice, C. W., Robledo Abad, C., Romanovskaya, A.,
30 Sperling, F., Herrero, M., House, J. I. and Rose, S.: How much land-based greenhouse gas mitigation
31 can be achieved without compromising food security and environmental goals?, *Glob. Change Biol.*,
32 19(8), 2285–2302, doi:10.1111/gcb.12160, 2013.
- 33 Stehfest, E., van Vuuren, D., Bouwman, L. and Kram, T.: *Integrated assessment of global environmental
34 change with IMAGE 3.0: Model description and policy applications.*, 2014.
- 35 van Straaten, O., Corre, M. D., Wolf, K., Tchienkoua, M., Cuellar, E., Matthews, R. B. and Veldkamp, E.:
36 Conversion of lowland tropical forests to tree cash crop plantations loses up to one-half of stored soil
37 organic carbon, *Proc. Natl. Acad. Sci.*, 112(32), 9956–9960, doi:10.1073/pnas.1504628112, 2015.
- 38 Vanninen, P., Ylitalo, H., Sievänen, R. and Mäkelä, A.: Effects of age and site quality on the distribution
39 of biomass in Scots pine (*Pinus sylvestris* L.), *Trees*, 10(4), 231–238, doi:10.1007/BF02185674, 1996.
- 40 Von Gadow, K. and Hui, G.: *Modelling Forest Development*, Springer Netherlands, Dordrecht., 1999.



- 1 van Vuuren, D. P., Deetman, S., van Vliet, J., van den Berg, M., van Ruijven, B. J. and Koelbl, B.: The role
- 2 of negative CO₂ emissions for reaching 2 °C—insights from integrated assessment modelling, *Clim.*
- 3 *Change*, 118(1), 15–27, doi:10.1007/s10584-012-0680-5, 2013.

- 4 Williamson, P.: Emissions reduction: Scrutinize CO₂ removal methods, *Nature*, 530(7589), 153–155,
- 5 doi:10.1038/530153a, 2016.

- 6 Zaehle, S., Sitch, S., Prentice, I. C., Liski, J., Cramer, W., Erhard, M., Hickler, T. and Smith, B.: The
- 7 Importance of Age-Related Decline in Forest NPP for Modeling Regional Carbon Balances, *Ecol. Appl.*,
- 8 16(4), 1555–1574, 2006.

- 9